

## PROPERTY ANALYSIS OF TRIPLE IMPLICATION METHOD FOR APPROXIMATE REASONING ON ATANASSOV'S INTUITIONISTIC FUZZY SETS

J. LI AND Y. LIU

**ABSTRACT.** Firstly, two kinds of natural distances between intuitionistic fuzzy sets are generated by the classical natural distance between fuzzy sets under a unified framework of residual intuitionistic implication operators. Secondly, the continuity and approximation property of a method for solving intuitionistic fuzzy reasoning are defined. It is proved that the triple implication method for intuitionistic fuzzy modus ponens has both continuity and approximation property with respect to these two kinds of natural distances based on Łukasiewicz implication, meanwhile the triple implication method for intuitionistic fuzzy modus tollens possesses unconditional continuity and conditional approximation property. Finally, some robustness results about the triple implication method of intuitionistic fuzzy reasoning are given.

### 1. Introduction

Since fuzzy set was firstly proposed by cybernetics expert Zadeh in 1965 (see [51]), it has provided a powerful tool to deal with fuzzy and uncertain information. It is well known that the fuzzy reasoning is a significant part of the theory of fuzzy sets. A flourishing achievement in fuzzy reasoning spreads out from the seminal paper of Zadeh [53] (see [8, 16, 18, 20, 22, 34, 35, 40, 43, 44, 46, 47, 48, 52, 54, 55]). Generally, two basic models of fuzzy reasoning are Fuzzy Modus Ponens and Fuzzy Modus Tollens. The basic reasoning principle underlying fuzzy modus ponens and Fuzzy Modus Tollens was Zadeh's Composition Rule of Inference (CRI). With the CRI method, fuzzy reasoning has been successfully applied to many fields (see [23, 32, 49]).

As an alternative for the CRI method, Wang [45] proposed a new method, called the full implication triple implication method, or simply the triple implication method, for fuzzy reasoning. This method may bring fuzzy reasoning within the framework of logical semantic implication, and it may be considered as a reasonable alternative or complement for the CRI method. Recently, many results of the triple implication method have been reported. Pei [36, 37] systematically discussed the method by using a class residual implications. Wang and Fu [42] gave the unified forms of the triple implication method by using the regular implication. Liu and Wang [29] obtained a class of restriction triple implication solution for

---

Received: xxxxxx; Revised: xxxxxx; Accepted: xxxxxx

*Key words and phrases:* Intuitionistic fuzzy reasoning, Triple implication method, Continuity, Approximation property, Robustness.

fuzzy reasoning. Pei [33] discussed the formalization problem of the triple implication fuzzy reasoning based on some first order logic systems and gave a sound logic deduce system for the unified triple implication method.

At the same time, the fuzzy control system requires the reasoning method to satisfy the approximation principle. Similar to approximation property, the continuity, perturbation and robustness can be used as a judge to decide whether the property of the existing reasoning method is good or bad. So analysis of properties of the fuzzy reasoning methods is a very active area of fuzzy research. Whether CRI method and triple implication method hold the continuity and approximation property was deeply discussed by Xu (see [50]), and how the approximation errors are propagated by them was given at last. Liu [27] carefully investigated the continuity and uniform continuity of the triple implication method. Dai discussed perturbation of fuzzy sets and fuzzy reasoning based on normalized Minkowski distances in [11].

While robustness analysis of fuzzy reasoning methods appears vital in particular. The class of t-norms and implications which are copulas and  $1-l_\infty$ -Lipschitz can guarantee good robustness of rule-based fuzzy reasoning (see [7]). In the literature [10], minimum similarity was used as a tool to analyse the robustness of triple implication method for fuzzy reasoning. However, Wang and Duan revised the concept of  $(T, \delta)$  equality as a measure to evaluate robustness of fuzzy reasoning and proved the continuity of the triple implication method with respect to the revised measures (see [41]). Li et al. [26] extended the concept of perturbations of fuzzy sets and studied the perturbations of fuzzy connectives and fuzzy reasoning based on the proposed divergence measures composed by DF-metric. He et al. [21] discussed the robustness of fuzzy connectives and fuzzy reasoning by using the new defined robustness index. Duan and Li [15] analyzed whether there are isolated points for the four specific logic metric spaces on  $F(X)$ , and they concluded that the logic metrics induced by Łukasiewicz implication and Goguen implication are more beneficial to the robustness analysis of fuzzy reasoning. From a knowledge representation point of view, fuzzy sets should be handled in some kinds of possible world semantics, however, the role of fuzziness is not always to capture uncertainty (see [17, 19]).

For instance, people make choices by checking the good sides and the bad sides of alternatives separately. Then they choose according to whether the good or the bad sides are stronger. The intuitionistic fuzzy sets introduced by Atanassov is a pair of fuzzy sets, namely a membership and a non-membership function (see [1, 2, 4, 5]), which can represent positive and negative aspects of the given information respectively, therefore it constitutes an appropriate knowledge representation framework, and has been widely applied in many fields such as pattern recognition, machine learning, decision making, image processing and so on (see [6, 12, 25, 28, 30]). But since Atanassovs intuitionistic fuzzy reasoning is not as natural as most of the fuzzy reasoning, especially restrictions on the intuitionistic fuzzy implication operators are much more complicated than that of fuzzy implication operators, so it seems that approximate reasoning based on Atanassovs intuitionistic fuzzy sets have not been given enough attention to.

In recent years, some scholars try to construct a logic foundation for intuitionistic fuzzy reasoning. Deschrijver et al. [13, 14] have made fruitful pioneering work, they presented the intuitionistic fuzzy t-norms and intuitionistic fuzzy implication operator theory. Begum and Srinivasan [39] defined the basic operations on the Intuitionistic Fuzzy Sets of Third Type (IFSTT) and established the relation between the modal operators like Necessity and Possibility. Cornelis and Deschrijver [9] put forward the ICRI method for intuitionistic fuzzy reasoning problems (i.e., IFMP and IFMT) by extending the CRI method to the intuitionistic fuzzy setting. Zheng et al. [56, 57] presented the representation of intuitionistic fuzzy implications by fuzzy operators, they extended the triple implication method of fuzzy reasoning on intuitionistic fuzzy reasoning and showed that the triple implication method of Intuitionistic Fuzzy Modus Ponens possessed virtue of reductivity and the triple implication method of Intuitionistic Fuzzy Modus Ponens was local reductive.

But so far, whether the existing triple implication method of intuitionistic fuzzy reasoning satisfies the approximation property? How about the continuity and robustness of it? These issues have not yet been studied. The present paper aims to investigate the properties of continuity, approximation and robustness of the triple implication method of intuitionistic fuzzy reasoning by extending the natural distance between fuzzy sets to intuitionistic fuzzy sets based on two basic inference models: Intuitionistic fuzzy modus ponens and Intuitionistic fuzzy modus tollens, and some results about continuity, approximation and robustness are proved based on Łukasiewicz implication.

The rest of this paper is organized as follows. In section 2, we review some necessary definitions, propositions and theorems, and propose the definition of continuity and approximation properties in the end. In section 3, we use some lemmas related to this paper to discuss the continuity, approximation properties and robustness of triple implication method for intuitionistic fuzzy reasoning with respect to Intuitionistic Fuzzy Modus Ponens and Intuitionistic Fuzzy Modus Tollens models. The final section includes our conclusions.

## 2. Preliminaries

In this section, we firstly review some basic concepts and results about intuitionistic fuzzy sets.

**Definition 2.1.** [1, 2, 4, 5] An intuitionistic fuzzy set on the nonempty universe of discourse  $X$  is given by  $A = \{(x, A_t(x), A_f(x)) | x \in X\}$  where

$$A_t : X \rightarrow [0, 1], A_f : X \rightarrow [0, 1]$$

with the condition

$$0 \leq A_t(x) + A_f(x) \leq 1, \forall x \in X.$$

$A_t(x)$  and  $A_f(x)$  denote the membership function and the non-membership function of  $x$  to  $A$ , respectively.

It is clear that the intuitionistic fuzzy set  $A$  in  $X$  can be written as:

$$A(x) = (u, v), 0 \leq u + v \leq 1, u, v \in [0, 1], \forall x \in X.$$

As a generalization of fuzzy sets, intuitionistic fuzzy sets extend the character value from  $[0, 1]$  to the triangle domain  $L^* = \{(u, v) \in [0, 1]^2 | u + v \leq 1\}$  (for a detailed geometrical interpretation of  $L^*$ , please see [1, 3, 5]).

If  $A_t(x) = 1 - A_f(x)$ ,  $\forall x \in X$ , then intuitionistic fuzzy sets degenerate into fuzzy sets. We denote  $\text{IF}(X)$  as the set of all intuitionistic fuzzy sets in  $X$  and  $F(X)$  as the set of all fuzzy sets in  $X$ .

Let  $\alpha, \beta \in L^*$ ,  $\alpha = (a_1, a_2)$ ,  $\beta = (b_1, b_2)$ , we can define a partial order on  $L^*$  as follows:

$$\alpha \leq \beta \text{ iff } a_1 \leq b_1, a_2 \geq b_2.$$

Obviously,  $\alpha \wedge \beta = (a_1 \wedge b_1, a_2 \vee b_2)$ ,  $\alpha \vee \beta = (a_1 \vee b_1, a_2 \wedge b_2)$ ,  $0^* = (0, 1)$  and  $1^* = (1, 0)$  are the smallest element and the greatest element of  $L^*$ , respectively. It's easy to verify the fact that  $(L^*, \leq)$  is a complete lattice.

**Definition 2.2.** [8, 13, 14] (i) A function  $\otimes_{L^*} : L^{*2} \rightarrow L^*$  is an intuitionistic fuzzy triangular norm (t-norm in short), if it is a commutative, associative and increasing function with neutral element  $1^*$ .

(ii) A function  $\oplus_{L^*} : L^{*2} \rightarrow L^*$  is an intuitionistic fuzzy triangular conorm (t-conorm in short), if it is a commutative, associative and increasing function with neutral element  $0^*$ .

(iii) A binary function  $\rightarrow_{L^*} : L^{*2} \rightarrow L^*$  satisfying the conditions:

$$0^* \rightarrow_{L^*} 0^* = 0^* \rightarrow_{L^*} 1^* = 1^* \rightarrow_{L^*} 1^* = 1^*$$

and

$$1^* \rightarrow_{L^*} 0^* = 0^*$$

is called an intuitionistic fuzzy implication.

(iv) An intuitionistic fuzzy negation  $' : L^* \rightarrow L^*$  satisfies, for all  $\alpha, \beta \in L^*$ , the following properties:

- (1)  $0^{*'} = 1^*$  and  $1^{*' } = 0^*$ .
- (2) If  $\alpha \geq \beta$ , then  $\alpha' \leq \beta'$ .

Now we give two binary operations  $\otimes_{L^*}, \oplus_{L^*}$  on  $L^*$  as follows:

$$\alpha \otimes_{L^*} \beta = (a_1 \otimes b_1, a_2 \oplus b_2), \quad \alpha \oplus_{L^*} \beta = (a_1 \oplus b_1, a_2 \otimes b_2),$$

where  $\oplus$  is the dual t-conorm of the t-norm  $\otimes$ .

First, we should clarify that two binary operations are effective, i.e.,  $\alpha \otimes_{L^*} \beta, \alpha \oplus_{L^*} \beta \in L^*$ . Since  $\alpha, \beta \in L^*$ , i.e.,  $(a_1, a_2) \in [0, 1]^2$ ,  $(b_1, b_2) \in [0, 1]^2$ ,  $a_1 + a_2 \leq 1$ ,  $b_1 + b_2 \leq 1$ , it follows from the monotonicity of  $\oplus$  and the duality of  $\otimes, \oplus$  that  $a_1 \otimes b_1 + a_2 \oplus b_2 \leq a_1 \otimes b_1 + (1 - a_1) \oplus (1 - b_1) = a_1 \otimes b_1 + 1 - a_1 \otimes b_1 = 1$ , i.e.,  $\alpha \otimes_{L^*} \beta \in L^*$ .

Similarly,  $\alpha \oplus_{L^*} \beta \in L^*$ .

It is easy to verify that  $\otimes_{L^*}$  and  $\oplus_{L^*}$  as defined above are intuitionistic t-norm and intuitionistic t-conorm respectively.

**Definition 2.3.** [31] Let  $X$  be non-empty universe of discourse,  $A_1, A_2 \in F(X)$ ,  $F(X)$  denotes the set of all fuzzy sets in  $X$ , then the natural distance between  $A_1$

and  $A_2$  is given by:

$$d_n(A_1, A_2) = \bigvee_{x \in X} |A_1(x) - A_2(x)|.$$

**Definition 2.4.** Let  $X$  be non-empty universe of discourse,  $A_1, A_2 \in IF(X)$ , then the natural distance between  $A_1$  and  $A_2$  can be given by:

- (i)  $d_{N_1}(A_1, A_2) = d_n(A_{1t}, A_{2t}) \vee d_n(A_{1f}, A_{2f})$ .
- (ii)  $d_{N_2}(A_1, A_2) = \frac{1}{2} \left( d_n(A_{1t}, A_{2t}) + d_n(A_{1f}, A_{2f}) \right)$ .

*Proof.* We only prove that  $d_{N_1}$  is a distance on  $IF(X)$ , the proof of  $d_{N_2}$  can be given in a similar way.

Firstly, for any  $A_1 = (A_{1t}, A_{1f}), A_2 = (A_{2t}, A_{2f}) \in IF(X)$ , it is clear that  $d_{N_1}(A_1, A_2) \geq 0$ . Suppose that  $A_1 = A_2$ , then we have  $A_{1t} = A_{2t}$  and  $A_{1f} = A_{2f}$ . Notice that  $d_n$  is the natural distance on  $F(X)$ , so it follows from  $d_n(A_{1t}, A_{2t}) = d_n(A_{1f}, A_{2f}) = 0$  that  $d_{N_1}(A_1, A_2) = 0$ . Conversely, if  $d_{N_1}(A_1, A_2) = 0$ , i.e.,  $d_n(A_{1t}, A_{2t}) \vee d_n(A_{1f}, A_{2f}) = 0$ , then  $d_n(A_{1t}, A_{2t}) = d_n(A_{1f}, A_{2f}) = 0$ , we have  $A_{1t} = A_{2t}$  and  $A_{1f} = A_{2f}$ . Therefore,  $A_1 = A_2$ .

Secondly, it is obvious that  $d_{N_1}(A_1, A_2) = d_{N_1}(A_2, A_1)$  holds for any  $A_1, A_2 \in IF(X)$ .

Thirdly, let  $A_1 = (A_{1t}, A_{1f}), A_2 = (A_{2t}, A_{2f}), A_3 = (A_{3t}, A_{3f}) \in IF(X)$ , it follows from the transitivity of  $d_n$  that  $d_n(A_{1t}, A_{3t}) \leq d_n(A_{1t}, A_{2t}) + d_n(A_{2t}, A_{3t})$  and  $d_n(A_{1f}, A_{3f}) \leq d_n(A_{1f}, A_{2f}) + d_n(A_{2f}, A_{3f})$ . Since

$$\begin{aligned} d_{N_1}(A_1, A_3) &= d_n(A_{1t}, A_{3t}) \vee d_n(A_{1f}, A_{3f}) \\ &\leq [d_n(A_{1t}, A_{2t}) + d_n(A_{2t}, A_{3t})] \vee [d_n(A_{1f}, A_{2f}) + d_n(A_{2f}, A_{3f})] \\ &\leq [d_n(A_{1t}, A_{2t}) \vee d_n(A_{1f}, A_{2f})] + [d_n(A_{2t}, A_{3t}) \vee d_n(A_{2f}, A_{3f})] \\ &\leq d_{N_1}(A_1, A_2) + d_{N_1}(A_2, A_3). \end{aligned}$$

In summary,  $d_{N_1}$  is a distance on  $IF(X)$ . This completes the proof.  $\square$

**Remark 2.5.** It can be easily proved that both  $d_{N_1}$  and  $d_{N_2}$  given by definition 2.3 are distances on  $IF(X)$ .

**Proposition 2.6.** [56, 57]  $(L^*, \otimes_{L^*}, 1^*)$  is a commutative monoid and  $\otimes_{L^*}$  is isotone,  $(L^*, \oplus_{L^*}, 0^*)$  is a commutative monoid and  $\oplus_{L^*}$  is isotone.

**Proposition 2.7.** [56, 57] Let  $\otimes$  be a left-continuous t-norm and  $I$  be the index set, then

- (i)  $\otimes_{L^*}$  is a left-continuous intuitionistic t-norm on  $L^*$ , i.e.,

$$\left( \bigvee_{i \in I} \alpha_i \right) \otimes_{L^*} \gamma = \bigvee_{i \in I} (\alpha_i \otimes_{L^*} \gamma), \forall \alpha_i, \gamma \in L^*.$$

- (ii)  $\oplus_{L^*}$  is a right-continuous intuitionistic t-conorm on  $L^*$ , i.e.,

$$\left( \bigwedge_{i \in I} \alpha_i \right) \oplus_{L^*} \gamma = \bigwedge_{i \in I} (\alpha_i \oplus_{L^*} \gamma), \forall \alpha_i, \gamma \in L^*.$$

**Theorem 2.8.** [56, 57] *Let  $\otimes_{L^*}$  be an intuitionistic t-norm induced by a left-continuous t-norm  $\otimes$ , then there exists a binary operation  $\rightarrow_{L^*}$  on  $L^*$  such that*

$$\gamma \otimes_{L^*} \alpha \leq \beta \text{ iff } \gamma \leq \alpha \rightarrow_{L^*} \beta, \quad (1)$$

and  $\rightarrow_{L^*}$  is given by

$$\alpha \rightarrow_{L^*} \beta = \bigvee \{ \eta \in L^* \mid \eta \otimes_{L^*} \alpha \leq \beta \}. \quad (2)$$

**Definition 2.9.** [56]  $(\otimes_{L^*}, \rightarrow_{L^*})$  is called an intuitionistic adjoint pair and  $\rightarrow_{L^*}$  is called a residual intuitionistic implication if  $(\otimes_{L^*}, \rightarrow_{L^*})$  satisfies the residual principle (1).

**Proposition 2.10.** [56] *Let  $\otimes_{L^*}$  be an intuitionistic t-norm on  $L^*$  and  $(\otimes_{L^*}, \rightarrow_{L^*})$  be an intuitionistic adjoint pair on  $L^*$ , then*

- (i)  $\alpha \rightarrow_{L^*} \beta = 1^*$  iff  $\alpha \leq \beta$ .
- (ii)  $\gamma \leq \alpha \rightarrow_{L^*} \beta$  iff  $\alpha \leq \gamma \rightarrow_{L^*} \beta$ .
- (iii)  $\gamma \rightarrow_{L^*} (\alpha \rightarrow_{L^*} \beta) = \alpha \rightarrow_{L^*} (\gamma \rightarrow_{L^*} \beta)$ .
- (iv)  $1^* \rightarrow_{L^*} \alpha = \alpha$ .
- (v)  $\beta \rightarrow_{L^*} \bigwedge_{i \in I} \alpha_i = \bigwedge_{i \in I} (\beta \rightarrow_{L^*} \alpha_i)$ .
- (vi)  $\bigvee_{i \in I} \beta_i \rightarrow_{L^*} \alpha = \bigwedge_{i \in I} (\beta_i \rightarrow_{L^*} \alpha)$ .
- (vii)  $\rightarrow_{L^*}$  is antitone in the first variable and isotone in the second variable.

Now we extend the triple implication method of Fuzzy Modus Ponens and Fuzzy Modus Tollens on intuitionistic fuzzy sets. We consider the problem of Intuitionistic Fuzzy Modus Ponens and Intuitionistic Fuzzy Modus Tollens as follows:

Suppose that	$A(x) \rightarrow_{L^*} B(y)$	...	major premise
and given	$A^*(x)$	...	minor premise
calculate	$B^*(y)$	...	conclusion
Suppose that	$A(x) \rightarrow_{L^*} B(y)$	...	major premise
and given	$B^*(y)$	...	minor premise
calculate	$A^*(x)$	...	conclusion

where  $A(x), A^*(x)$  are the intuitionistic fuzzy sets on  $X$ , respectively,  $B(y), B^*(y)$  are the intuitionistic fuzzy sets on  $Y$ , respectively, and  $\rightarrow_{L^*}$  is the residual intuitionistic fuzzy implication on  $L^*$ . We denote  $A(x) = (A_t(x), A_f(x)), B(y) = (B_t(y), B_f(y)), A^*(x) = (A_t^*(x), A_f^*(x)), B^*(y) = (B_t^*(y), B_f^*(y)), A_{-f}(x) = 1 - A_f(x), B_{-f}(y) = 1 - B_f(y), A_{-f}^*(x) = 1 - A_f^*(x), B_{-f}^*(y) = 1 - B_f^*(y)$ . Clearly,  $A_t, A_f, A_t^*, A_f^*, A_{-f}, A_{-f}^*$  are the fuzzy sets on  $X$ , respectively, and  $B_t, B_f, B_t^*, B_f^*, B_{-f}, B_{-f}^*$  are the fuzzy sets on  $Y$ , respectively.

Because  $\rightarrow_{L^*}$  is the residual intuitionistic fuzzy implication on  $L^*$ , we could smoothly reach the extension of the Triple Implication Principle for Intuitionistic Fuzzy Modus Ponens as follows:  $B^*(y)$  should be the smallest intuitionistic fuzzy set on  $Y$  satisfying

$$(A(x) \rightarrow_{L^*} B(y)) \rightarrow_{L^*} (A^*(x) \rightarrow_{L^*} B^*(y)) = 1^*, \quad (3)$$

under the order of  $L^*$ .

Similarly, the Triple Implication Principle for Intuitionistic Fuzzy Modus Tollens can be extended as follows:  $A^*(x)$  should be the biggest intuitionistic fuzzy set on  $X$  satisfying (3) under the order of  $L^*$ .

**Theorem 2.11.** [56, 57] *Suppose that the implication  $\rightarrow_{L^*}$  in IFMP is the residual intuitionistic implication induced by a left-continuous  $t$ -norm  $\otimes$ , then the triple implication solution  $B^*$  of IFMP is given by the following formula*

$$B^*(y) = \bigvee_{x \in X} \{A^*(x) \otimes_{L^*} (A(x) \rightarrow_{L^*} B(y))\}, \quad y \in Y. \quad (4)$$

and the decomposition expression of  $B^*(y)$  is given by

$$B^*(y) = \left( \bigvee_{x \in X} \left\{ A_t^*(x) \otimes \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \right\}, \right. \\ \left. \bigwedge_{x \in X} \left\{ A_f^*(x) \oplus (1 - A_{-f}(x) \rightarrow B_{-f}(y)) \right\} \right). \quad (5)$$

**Theorem 2.12.** [56] *Let the implication  $\rightarrow_{L^*}$  in IFMT be the residual intuitionistic implication induced by a left-continuous  $t$ -norm  $\otimes$ , then the triple implication solution  $A^*$  of IFMT is given by the following formula*

$$A^*(x) = \bigvee_{y \in Y} \left\{ (A(x) \rightarrow_{L^*} B(y)) \rightarrow_{L^*} B^*(y) \right\}, \quad x \in X. \quad (6)$$

and the decomposition expression of  $A^*(x)$  is given by

$$A^*(x) = \left( \bigwedge_{y \in Y} \left\{ \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \rightarrow B_t^*(y) \right\}, \right. \\ \left. \bigwedge \left\{ (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}^*(y) \right\}, \right. \\ \left. \bigvee_{y \in Y} \left\{ 1 - (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}^*(y) \right\} \right). \quad (7)$$

**Definition 2.13.** Let  $X, Y$  be non-empty universe of discourse,  $A \rightarrow B$  be the known rule. A inference method for solving IFMP is an  $IF(X) \rightarrow IF(Y)$  mapping  $g$ , i.e., each input  $A^* \in IF(X)$  corresponds to an output  $B^* = g(A^*)$ . Let  $d_N$  be the natural distance between intuitionistic fuzzy sets.

- (i)  $g$  is said to be continuous at  $A$  under metric  $d_N$ , if for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $d_N(g(A^*), g(A)) < \varepsilon$  whenever  $d_N(A^*, A) < \delta$  for any  $A^* \in IF(X)$ .
- (ii)  $g$  is said to be of approximation property at  $B \in IF(Y)$  under metric  $d_N$ , if for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $d_N(g(A^*), B) < \varepsilon$  whenever  $d_N(A^*, A) < \delta$  for any  $A^* \in IF(X)$ .
- (iii)  $g$  is said to be reductive if  $g(A) = B$  holds, i.e.,  $B^* = B$  holds whenever  $A^* = A$ .

**Definition 2.14.** Let  $X, Y$  be non-empty universe of discourse,  $A \rightarrow B$  be the known rule. A method for solving IFMT is an  $IF(Y) \rightarrow IF(X)$  mapping  $h$ , i.e., each input  $B^* \in IF(Y)$  corresponds to an output  $A^* = h(B^*)$ . Let  $d_N$  be the natural distance between intuitionistic fuzzy sets.

- (i)  $h$  is said to be continuous at  $B$  under metric  $d_N$ , if for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $d_N(h(B^*), h(B)) < \varepsilon$  whenever  $d_N(B^*, B) < \delta$  for any  $B^* \in IF(Y)$ .
- (ii)  $h$  is said to be of approximation property at  $A \in IF(X)$  under metric  $d_N$ , if for any  $\varepsilon > 0$ , there exists  $\delta > 0$  such that  $d_N(h(B^*), A) < \varepsilon$  whenever  $d_N(B^*, B) < \delta$  for any  $B^* \in IF(Y)$ .
- (iii)  $h$  is said to be reductive if  $h(B) = A$  holds, i.e.,  $A^* = A$  holds whenever  $B^* = B$ .

**Remark 2.15.** The choice of distance in Definition 2.13 and Definition 2.14 is not unique. It could be any other one, such as the Hamming distance and the Euclidean distance.

**Remark 2.16.** (i) It can be easily seen that an inference method  $g$  for Intuitionistic Fuzzy Modus Ponens must be of approximation property if  $g$  is both continuous and reductive, and the same conclusion also holds with respect to the method  $h$  for Intuitionistic Fuzzy Modus Tollens.

(ii) The continuity of  $g$  reflects the fact that if the input  $A^*$  is very close to  $A$  then the output  $B^*$  (i.e.,  $g(A^*)$ ) can be very close to  $g(A)$ . The approximation property of  $g$  means that if the input  $A^*$  is very close to  $A$  then the output  $B^*$  is very close to  $B$ .

A similar analysis can be done with the method for Intuitionistic Fuzzy Modus Tollens.

### 3. The continuity, approximation property and robustness of triple implication method

As is well known that fuzzy logic is the mathematical foundation of fuzzy reasoning while the completeness theorem is one of the important criteria to judge whether logic system is good or bad, the fuzzy logic system based on Łukasiewicz implication is one of the most important fuzzy logic system due to the completeness theorem holds in it. In addition, since Łukasiewicz implication is continuous and is of good transitivity, so it is one of the most common used implication operators in fuzzy reasoning.

In this paper, we focus on the analysis of  $\otimes_{L^*}$  and its residuum  $\rightarrow_{L^*}$  induced by Łukasiewicz t-norm. In the subsequent sections, unless otherwise stated,  $\otimes, \oplus, \rightarrow$  always represents Łukasiewicz t-norm, Łukasiewicz t-conorm and Łukasiewicz implication operator. And  $a \otimes b = (a+b-1) \vee 0$ ,  $a \oplus b = (a+b) \wedge 1$ ,  $a \rightarrow b = (1-a+b) \wedge 1$ . To facilitate future disussion, it is necessary to introduce the following three lemmas.

**Lemma 3.1.** [35] *Let  $f$  and  $g$  be bounded, real valued functions on a set  $X$ . Then*



$$(i) \quad \left| \bigvee_{x \in X} f(x) - \bigvee_{x \in X} g(x) \right| \leq \bigvee_{x \in X} |f(x) - g(x)|,$$

$$(ii) \quad \left| \bigwedge_{x \in X} f(x) - \bigwedge_{x \in X} g(x) \right| \leq \bigvee_{x \in X} |f(x) - g(x)|.$$

**Lemma 3.2.** For any  $a, b, c, d \in [0, 1]$ , then

$$(i) \quad |a \otimes b - c \otimes d| \leq |a - c| + |b - d|, |a \otimes b - c \otimes b| \leq |a - c|.$$

$$(ii) \quad |a \otimes b - c \otimes d| \leq 2(|a - c| \vee |b - d|).$$

$$(iii) \quad |a \oplus b - c \oplus d| \leq |a - c| + |b - d|, |a \oplus b - c \oplus b| \leq |a - c|.$$

$$(iv) \quad |a \oplus b - c \oplus d| \leq 2(|a - c| \vee |b - d|).$$

*Proof.* Firstly, it is obvious that  $|a - c| + |b - d| \leq 2(|a - c| \vee |b - d|)$ .

We only need to prove that  $|a \otimes b - c \otimes d| \leq |a - c| + |b - d|$ ,  $|a \oplus b - c \oplus d| \leq |a - c| + |b - d|$ .

By using Lemma 3.1(i), we can easily get that

$$|a \otimes b - c \otimes d| = \left| (a + b - 1) \vee 0 - (c + d - 1) \vee 0 \right|$$

$$\leq |a - c + b - d| \leq |a - c| + |b - d|.$$

It follows from Lemma 3.1(ii) that

$$|a \oplus b - c \oplus d| = \left| (a + b) \wedge 1 - (c + d) \wedge 1 \right|$$

$$\leq |a - c + b - d| \leq |a - c| + |b - d|.$$

This completes the proof.  $\square$

**Lemma 3.3.** For any  $a, b, c, d \in [0, 1]$ , then

$$(i) \quad |a \rightarrow b - c \rightarrow d| \leq |a - c| + |b - d|, |a \rightarrow b - c \rightarrow b| \leq |a - c|.$$

$$(ii) \quad |a \rightarrow b - c \rightarrow d| \leq 2(|a - c| \vee |b - d|).$$

*Proof.* Similarly, by using Lemma 3.1(ii), we have

$$|a \rightarrow b - c \rightarrow d| = \left| (1 - a + b) \wedge 1 - (1 - c + d) \wedge 1 \right|$$

$$\leq |c - a + b - d| \leq |a - c| + |b - d|.$$

This completes the proof.  $\square$

**Remark 3.4.** Lemma 3.2 also holds when  $\otimes$  is Minimum t-norm or Product t-norm.

**Theorem 3.5.** The triple implication method  $g$  for IFMP based on the Lukasiewicz implication is continuous at  $A$  under the metric  $d_{N_1}$ .

*Proof.* It follows from Definition 2.4 that

$$d_{N_1}(g(A^*), g(A)) = d_n(g(A^*)_t, g(A)_t) \vee d_n(g(A^*)_f, g(A)_f).$$

(8)

From (8) and (5), we can first obtain that

$$\begin{aligned}
d_n(g(A^*)_t, g(A)_t) &= \bigvee_{y \in Y} |g(A^*)_t(y) - g(A)_t(y)| \\
&= \bigvee_{y \in Y} \left| \bigvee_{x \in X} \left\{ A_t^*(x) \otimes \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \right\} \right. \\
&\quad \left. - \bigvee_{x \in X} \left\{ A_t(x) \otimes \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \right\} \right|. \tag{9}
\end{aligned}$$

Then by Lemma 3.1(i) and Lemma 3.2(i), we have

$$\begin{aligned}
d_n(g(A^*)_t, g(A)_t) &\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left| A_t^*(x) \otimes \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \right. \\
&\quad \left. - A_t(x) \otimes \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \right| \\
&\leq \bigvee_{x \in X} |A_t^*(x) - A_t(x)|. \tag{10}
\end{aligned}$$

Similarly, by (8) and (5), we can easily get that

$$\begin{aligned}
d_n(g(A^*)_f, g(A)_f) &= \bigvee_{y \in Y} |g(A^*)_f(y) - g(A)_f(y)| \\
&= \bigvee_{y \in Y} \left| \bigwedge_{x \in X} \left\{ A_f^*(x) \oplus (1 - A_{-f}(x) \rightarrow B_{-f}(y)) \right\} \right. \\
&\quad \left. - \bigwedge_{x \in X} \left\{ A_f(x) \oplus (1 - A_{-f}(x) \rightarrow B_{-f}(y)) \right\} \right|. \tag{11}
\end{aligned}$$

According to eq.(11), Lemma 3.1(ii) and Lemma 3.2(iii), it is obvious that

$$\begin{aligned}
d_n(g(A^*)_f, g(A)_f) &\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left| A_f^*(x) \oplus (1 - A_{-f}(x) \rightarrow B_{-f}(y)) \right. \\
&\quad \left. - A_f(x) \oplus (1 - A_{-f}(x) \rightarrow B_{-f}(y)) \right| \\
&\leq \bigvee_{y \in Y} \bigvee_{x \in X} |A_f^*(x) - A_f(x)| = \bigvee_{x \in X} |A_f^*(x) - A_f(x)|. \tag{12}
\end{aligned}$$

Therefore, for any  $\varepsilon > 0$ , by taking  $\delta = \varepsilon$ , when  $d_{N_1}(A^*, A) < \delta$ , then by eq.(8), (10) and (12) we have

$$\begin{aligned}
d_{N_1}(g(A^*), g(A)) &\leq \left( \bigvee_{x \in X} |A_t^*(x) - A_t(x)| \right) \bigvee \left( \bigvee_{x \in X} |A_f^*(x) - A_f(x)| \right) \\
&= d_{N_1}(A^*, A) < \delta = \varepsilon.
\end{aligned}$$

This completes the proof of Theorem.  $\square$

**Theorem 3.6.** *Based on the Lukasiewicz implication, the triple implication method  $g$  for IFMP is continuous at  $A$  under the metric  $d_{N_2}$ .*

*Proof.* It follows from Definition 2.4 that

$$d_{N_2}(g(A^*), g(A)) = \frac{1}{2} \left( d_n(g(A^*)_t, g(A)_t) + d_n(g(A^*)_f, g(A)_f) \right). \quad (13)$$

By using the same method as the proof of Theorem 3.5, we can obtain that

$$d_n(g(A^*)_t, g(A)_t) \leq \bigvee_{x \in X} |A_t^*(x) - A_t(x)|. \quad (14)$$

$$d_n(g(A^*)_f, g(A)_f) \leq \bigvee_{x \in X} |A_f^*(x) - A_f(x)|. \quad (15)$$

Therefore, for any  $\varepsilon > 0$ , by taking  $\delta = \varepsilon$ , when  $d_{N_1}(A^*, A) \leq \delta$ , then by eq.(13), (14) and (15) we have

$$\begin{aligned} d_{N_2}(g(A^*), g(A)) &\leq \frac{1}{2} \left( d_n(g(A^*)_t, g(A)_t) + d_n(g(A^*)_f, g(A)_f) \right) \\ &= d_{N_2}(A^*, A) < \delta = \varepsilon. \end{aligned}$$

This completes the proof of Theorem.  $\square$

**Lemma 3.7.** [56] *If  $\rightarrow_{L^*}$  is a residual intuitionistic implication induced by a left-continuous  $t$ -norm, then the triple implication method of IFMP given by Theorem 2.11 is reductive whenever  $A$  is a normal intuitionistic fuzzy set on  $X$ , i.e., there exists  $x_0$  in  $X$  such that  $A(x_0) = 1^*$ .*

The following Corollary can be easily get by Lemma 3.7 and Remark 2.16.

**Corollary 3.8.** *Let  $A \rightarrow B$  be the known rule,  $A \in IF(X)$ ,  $B \in IF(Y)$ . If  $A^*$  is a normal intuitionistic fuzzy set, then the triple implication method  $g$  for IFMP is of approximation property.*

**Theorem 3.9.** *The triple implication method  $h$  for IFMT based on the Lukasiewicz implication is continuous at  $B$  under the metric  $d_{N_1}$ .*

*Proof.* Firstly, it is easy for us to know that  $|B_{-f}^*(y) - B_{-f}(y)| = |B_f^*(y) - B_f(y)|$ . From eq.(7) and eq.(8), we could obtain that

$$\begin{aligned} d_n(h(B^*)_t, h(B)_t) &= \bigvee_{x \in X} |h(B^*)_t(x) - h(B)_t(x)| \\ &= \bigvee_{x \in X} \left| \bigwedge_{y \in Y} \left\{ \left( (A_t(x) \rightarrow B_t(y)) \bigwedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \rightarrow B_t^*(y) \right\} \right. \\ &\quad \left. \bigwedge \left( (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}^*(y) \right) \right\} \\ &\quad - \bigwedge_{y \in Y} \left\{ \left( (A_t(x) \rightarrow B_t(y)) \bigwedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \rightarrow B_t(y) \right\} \\ &\quad \left. \bigwedge \left( (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}(y) \right) \right\} \Big|. \end{aligned} \quad (16)$$

Then by using Lemma 3.1(ii) twice, we have

$$\begin{aligned}
& d_n(h(B^*)_t, h(B)_t) \\
& \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( \left| \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \rightarrow B_t^*(y) \right. \right. \\
& \quad \left. \left. - \left( (A_t(x) \rightarrow B_t(y)) \wedge (A_{-f}(x) \rightarrow B_{-f}(y)) \right) \rightarrow B_t(y) \right| \right) \\
& \quad \bigvee \left| (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}^*(y) - (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}(y) \right|. \quad (17)
\end{aligned}$$

It follows from Lemma 3.3(i) that

$$\begin{aligned}
d_n(h(B^*)_t, h(B)_t) & \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( |B_t^*(y) - B_t(y)| \bigvee |B_{-f}^*(y) - B_{-f}(y)| \right) \\
& = \bigvee_{y \in Y} \left( |B_t^*(y) - B_t(y)| \bigvee |B_{-f}^*(y) - B_{-f}(y)| \right) \\
& = \left( \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| \right) \bigvee \left( \bigvee_{y \in Y} |B_{-f}^*(y) - B_{-f}(y)| \right). \quad (18)
\end{aligned}$$

Similarly, by using eq.(7) and eq.(8), we can easily get that

$$\begin{aligned}
d_n(h(B^*)_f, h(B)_f) & = \bigvee_{x \in X} |h(B^*)_f(x) - h(B)_f(x)| \\
& = \bigvee_{x \in X} \left| \bigvee_{y \in Y} \{1 - (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}^*(y)\} \right. \\
& \quad \left. - \bigvee_{y \in Y} \{1 - (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}(y)\} \right|. \quad (19)
\end{aligned}$$

From (19), by using Lemma 3.1(i) and Lemma 3.3(i), it is obvious that

$$\begin{aligned}
d_n(h(B^*)_f, h(B)_f) & \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left| (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}^*(y) \right. \\
& \quad \left. - (A_{-f}(x) \rightarrow B_{-f}(y)) \rightarrow B_{-f}(y) \right| \\
& \leq \bigvee_{x \in X} \bigvee_{y \in Y} |B_{-f}^*(y) - B_{-f}(y)| = \bigvee_{y \in Y} |B_{-f}^*(y) - B_{-f}(y)|. \quad (20)
\end{aligned}$$

Therefore, for any  $\varepsilon > 0$ , by taking  $\delta = \varepsilon$ , when  $d_{N_1}(B^*, B) \leq \delta$ , then by eq.(8), (18) and (20), we can easily obtain that

$$\begin{aligned}
d_{N_1}(h(B^*), h(B)) & \leq \left( \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| \right) \bigvee \left( \bigvee_{y \in Y} |B_{-f}^*(y) - B_{-f}(y)| \right) \\
& \quad \bigvee \left( \bigvee_{y \in Y} |B_{-f}^*(y) - B_{-f}(y)| \right) \\
& = \left( \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| \right) \bigvee \left( \bigvee_{y \in Y} |B_{-f}^*(y) - B_{-f}(y)| \right) \\
& = d_{N_1}(B^*, B) < \delta = \varepsilon.
\end{aligned}$$

This completes the proof of Theorem.  $\square$

**Theorem 3.10.** *Based on the Lukasiewicz implication, the triple implication method  $h$  for IFMT is continuous at  $B$  under the metric  $d_{N_2}$ .*

*Proof.* Using the method same as the Theorem 3.9, we can obtain that

$$\begin{aligned} d_n(h(B^*)_t, h(B)_t) &\leq \left( \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| \right) \bigvee \left( \bigvee_{y \in Y} |B_f^*(y) - B_f(y)| \right) \\ &\leq \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| + \bigvee_{y \in Y} |B_f^*(y) - B_f(y)|. \end{aligned} \quad (21)$$

$$d_n(h(B^*)_f, h(B)_f) \leq \bigvee_{y \in Y} |B_f^*(y) - B_f(y)|. \quad (22)$$

Therefore, for any  $\varepsilon > 0$ , by taking  $\delta = \frac{1}{2}\varepsilon$ , when  $d_{N_1}(B^*, B) < \delta$ , then by eq.(13), (21) and (22), it is obvious that

$$\begin{aligned} d_{N_2}(h(B^*), h(B)) &\leq \frac{1}{2} \left( \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| + 2 \bigvee_{y \in Y} |B_f^*(y) - B_f(y)| \right) \\ &\leq \bigvee_{y \in Y} |B_t^*(y) - B_t(y)| + \bigvee_{y \in Y} |B_f^*(y) - B_f(y)| \\ &= 2d_{N_2}(B^*, B) \\ &< 2\delta = \varepsilon. \end{aligned}$$

Hence, there exists  $\delta > 0$  such that  $d_{N_2}(h(B^*), h(B)) < \varepsilon$  when  $d_{N_2}(B^*, B) < \delta$ . This completes the proof of the Theorem.  $\square$

**Theorem 3.11.** *Let  $B_1^*$  and  $B_2^*$  be the triple implication solutions of IFMP  $(A_1, B_1, A_1^*)$  and IFMP  $(A_2, B_2, A_2^*)$  based on the Lukasiewicz implication, respectively. Suppose that  $d_{N_1}(A_1, A_2) \leq \delta_1$ ,  $d_{N_1}(B_1, B_2) \leq \delta_2$ ,  $d_{N_1}(A_1^*, A_2^*) \leq \delta_3$ , then*

$$d_{N_1}(B_1^*, B_2^*) \leq 4\delta_1 \bigvee 4\delta_2 \bigvee 2\delta_3.$$

*Proof.* We firstly take the natural distance between the membership functions of  $B_1^*$  and  $B_2^*$  into consideration, by definition 2.3 and eq.(5) we have

$$\begin{aligned} &d_n(B_{1t}^*, B_{2t}^*) \\ &= \bigvee_{y \in Y} \left| \bigvee_{x \in X} \left\{ A_{1t}^*(x) \otimes \left( (A_{1t}(x) \rightarrow B_{1t}(y)) \bigwedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right) \right\} \right. \\ &\quad \left. - \bigvee_{x \in X} \left\{ A_{2t}^*(x) \otimes \left( (A_{2t}(x) \rightarrow B_{2t}(y)) \bigwedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right) \right\} \right|. \end{aligned}$$

(23)

From (23), Lemma 3.1(i) and Lemma 3.2(ii), we can easily obtain that

$$\begin{aligned}
& d_n(B_{1t}^*, B_{2t}^*) \\
& \leq \bigvee_{y \in Y} \bigvee_{x \in X} \left| A_{1t}^*(x) \otimes \left( (A_{1t}(x) \rightarrow B_{1t}(y)) \wedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right) \right. \\
& \quad \left. - A_{2t}^*(x) \otimes \left( (A_{2t}(x) \rightarrow B_{2t}(y)) \wedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right) \right| \\
& \leq \bigvee_{y \in Y} \bigvee_{x \in X} 2 \left( \left| A_{1t}^*(x) - A_{2t}^*(x) \right| \bigvee \left| (A_{1t}(x) \rightarrow B_{1t}(y)) \wedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right. \right. \\
& \quad \left. \left. - (A_{2t}(x) \rightarrow B_{2t}(y)) \wedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right| \right). \tag{24}
\end{aligned}$$

Then it follows from (24) and Lemma 3.1(ii) that

$$\begin{aligned}
& d_n(B_{1t}^*, B_{2t}^*) \\
& \leq \bigvee_{y \in Y} \bigvee_{x \in X} 2 \left( \left| A_{1t}^*(x) - A_{2t}^*(x) \right| \bigvee \left| A_{1t}(x) \rightarrow B_{1t}(y) - A_{2t}(x) \rightarrow B_{2t}(y) \right| \right. \\
& \quad \left. \bigvee \left| A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y) \right| \right). \tag{25}
\end{aligned}$$

According to Lemma 3.3 (ii), we further have

$$\begin{aligned}
& d_n(B_{1t}^*, B_{2t}^*) \leq \bigvee_{y \in Y} \bigvee_{x \in X} 2 \left( \left| A_{1t}^*(x) - A_{2t}^*(x) \right| \bigvee 2 \left( \left| A_{1t}(x) - A_{2t}(x) \right| \bigvee \left| B_{1t}(y) \right. \right. \right. \\
& \quad \left. \left. - B_{2t}(y) \right| \right) \bigvee 2 \left( \left| A_{1-f}(x) - A_{2-f}(x) \right| \bigvee \left| B_{1-f}(y) - B_{2-f}(y) \right| \right) \\
& = \left( 2 \bigvee_{x \in X} \left| A_{1t}^*(x) - A_{2t}^*(x) \right| \right) \bigvee \left( 4 \bigvee_{x \in X} \left| A_{1t}(x) - A_{2t}(x) \right| \right) \\
& \quad \bigvee \left( 4 \bigvee_{y \in Y} \left| B_{1t}(y) - B_{2t}(y) \right| \right) \bigvee \left( 4 \bigvee_{x \in X} \left| A_{1f}(x) - A_{2f}(x) \right| \right) \\
& \quad \bigvee \left( 4 \bigvee_{y \in Y} \left| B_{1f}(y) - B_{2f}(y) \right| \right). \tag{26}
\end{aligned}$$

Secondly, we consider the natural distance between the non-membership functions of  $B_{1f}^*$  and  $B_{2f}^*$ , by Definition 2.3, eq.(5) and Lemma 3.1(ii) we have

$$\begin{aligned}
& d_n(B_{1f}^*, B_{2f}^*) = \bigvee_{y \in Y} \left| \bigwedge_{x \in X} \left\{ A_{1f}^*(x) \oplus (1 - A_{1-f}(x) \rightarrow B_{1-f}(y)) \right\} \right. \\
& \quad \left. - \bigwedge_{x \in X} \left\{ A_{2f}^*(x) \oplus (1 - A_{2-f}(x) \rightarrow B_{2-f}(y)) \right\} \right| \\
& \leq \bigvee_{y \in Y} \bigvee_{x \in X} \left| A_{1f}^*(x) \oplus (1 - A_{1-f}(x) \rightarrow B_{1-f}(y)) \right. \\
& \quad \left. - A_{2f}^*(x) \oplus (1 - A_{2-f}(x) \rightarrow B_{2-f}(y)) \right|. \tag{27}
\end{aligned}$$

Then it can be easily gotten from (27), Lemma 3.2(iv) and Lemma 3.3(ii) that

$$\begin{aligned}
d_n(B_{1f}^*, B_{2f}^*) &\leq \bigvee_{y \in Y} \bigvee_{x \in X} 2 \left( |A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y)| \right. \\
&\quad \left. \bigvee |A_{1f}^*(x) - A_{2f}^*(x)| \right) \\
&\leq \left( 2 \bigvee_{x \in X} |A_{1t}^*(x) - A_{2t}^*(x)| \right) \bigvee \left( 4 \bigvee_{x \in X} |A_{1f}(x) - A_{2f}(x)| \right) \\
&\quad \bigvee \left( 4 \bigvee_{y \in Y} |B_{1f}(y) - B_{2f}(y)| \right).
\end{aligned} \tag{28}$$

Thus it follows from eq.(8), (26) and (28) that

$$\begin{aligned}
d_{N_1}(B_1^*, B_2^*) &\leq \left( 2 \bigvee_{x \in X} |A_{1t}^*(x) - A_{2t}^*(x)| \right) \bigvee \left( 2 \bigvee_{x \in X} |A_{1f}^*(x) - A_{2f}^*(x)| \right) \\
&\quad \bigvee \left( 4 \bigvee_{x \in X} |A_{1t}(x) - A_{2t}(x)| \right) \bigvee \left( 4 \bigvee_{x \in X} |A_{1f}(x) - A_{2f}(x)| \right) \\
&\quad \bigvee \left( 4 \bigvee_{y \in Y} |B_{1t}(y) - B_{2t}(y)| \right) \bigvee \left( 4 \bigvee_{y \in Y} |B_{1f}(y) - B_{2f}(y)| \right) \\
&= 4d_{N_1}(A_1, A_2) \bigvee 4d_{N_1}(B_1, B_2) \bigvee 2d_{N_1}(A_1^*, A_2^*) \\
&\leq 4\delta_1 \bigvee 4\delta_2 \bigvee 2\delta_3.
\end{aligned}$$

This completes the proof.  $\square$

**Theorem 3.12.** *Let  $B_1^*$  and  $B_2^*$  be the triple implication solutions of IFMP  $(A_1, B_1, A_1^*)$  and IFMP  $(A_2, B_2, A_2^*)$  based on the Łukasiewicz implication, respectively. Suppose that  $d_{N_2}(A_1, A_2) \leq \delta_1$ ,  $d_{N_2}(B_1, B_2) \leq \delta_2$ ,  $d_{N_2}(A_1^*, A_2^*) \leq \delta_3$ , then*

$$d_{N_2}(B_1^*, B_2^*) \leq 2\delta_1 + 2\delta_2 + \delta_3.$$

*Proof.* Firstly we take the natural distance between the membership functions of  $B_1^*$  and  $B_2^*$  into consideration, by definition 2.3 and eq.(5) we have

$$\begin{aligned}
d_n(B_{1t}^*, B_{2t}^*) &= \bigvee_{y \in Y} \left| \bigvee_{x \in X} \left\{ A_{1t}^*(x) \otimes \left( (A_{1t}(x) \rightarrow B_{1t}(y)) \wedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right) \right\} \right. \\
&\quad \left. - \bigvee_{x \in X} \left\{ A_{2t}^*(x) \otimes \left( (A_{2t}(x) \rightarrow B_{2t}(y)) \wedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right) \right\} \right|.
\end{aligned}$$

(29)

From (29), Lemma 3.1(i) and Lemma 3.2(i), it can be easily obtained that

$$\begin{aligned}
d_n(B_{1t}^*, B_{2t}^*) &\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left| A_{1t}^*(x) \otimes \left( (A_{1t}(x) \rightarrow B_{1t}(y)) \bigwedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right) \right. \\
&\quad \left. - A_{2t}^*(x) \otimes \left( (A_{2t}(x) \rightarrow B_{2t}(y)) \bigwedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right) \right| \\
&\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left( |A_{1t}^*(x) - A_{2t}^*(x)| + \left| (A_{1t}(x) \rightarrow B_{1t}(y)) \bigwedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right. \right. \\
&\quad \left. \left. - (A_{2t}(x) \rightarrow B_{2t}(y)) \bigwedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right| \right). \tag{30}
\end{aligned}$$

Then it follows from (30) and Lemma 3.1(ii) that

$$\begin{aligned}
d_n(B_{1t}^*, B_{2t}^*) &\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left( |A_{1t}^*(x) - A_{2t}^*(x)| + |A_{1t}(x) \rightarrow B_{1t}(y) - A_{2t}(x) \rightarrow B_{2t}(y)| \right. \\
&\quad \left. \bigvee |A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y)| \right) \\
&\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left( |A_{1t}^*(x) - A_{2t}^*(x)| + |A_{1t}(x) \rightarrow B_{1t}(y) - A_{2t}(x) \rightarrow B_{2t}(y)| \right. \\
&\quad \left. + |A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y)| \right). \tag{31}
\end{aligned}$$

For (31), according to Lemma 3.3(i), we further have

$$\begin{aligned}
d_n(B_{1t}^*, B_{2t}^*) &\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left( |A_{1t}^*(x) - A_{2t}^*(x)| + |A_{1t}(x) - A_{2t}(x)| + |B_{1t}(y) - B_{2t}(y)| \right. \\
&\quad \left. + |A_{1-f}(x) - A_{2-f}(x)| + |B_{1-f}(y) - B_{2-f}(y)| \right) \\
&\leq \bigvee_{x \in X} |A_{1t}^*(x) - A_{2t}^*(x)| + \bigvee_{x \in X} |A_{1t}(x) - A_{2t}(x)| + \bigvee_{y \in Y} |B_{1t}(y) - B_{2t}(y)| \\
&\quad + \bigvee_{x \in X} |A_{1-f}(x) - A_{2-f}(x)| + \bigvee_{y \in Y} |B_{1-f}(y) - B_{2-f}(y)|. \tag{32}
\end{aligned}$$

Secondly, we consider the natural distance between the non-membership functions of  $B_1^*$  and  $B_2^*$ , by Definition 2.3, eq.(5) and Lemma 3.1(ii) we have

$$\begin{aligned}
d_n(B_{1f}^*, B_{2f}^*) &= \bigvee_{y \in Y} \left| \bigwedge_{x \in X} \{ A_{1f}^*(x) \oplus (1 - A_{1-f}(x) \rightarrow B_{1-f}(y)) \} \right. \\
&\quad \left. - \bigwedge_{x \in X} \{ A_{2f}^*(x) \oplus (1 - A_{2-f}(x) \rightarrow B_{2-f}(y)) \} \right| \\
&\leq \bigvee_{y \in Y} \bigvee_{x \in X} \left| A_{1f}^*(x) \oplus (1 - A_{1-f}(x) \rightarrow B_{1-f}(y)) \right. \\
&\quad \left. - A_{2f}^*(x) \oplus (1 - A_{2-f}(x) \rightarrow B_{2-f}(y)) \right|. \tag{33}
\end{aligned}$$



Then it can be easily gotten from (33), Lemma 3.2(iii) and Lemma 3.3(i) that

$$\begin{aligned}
& d_n(B_{1f}^*, B_{2f}^*) \\
& \leq \bigvee_{y \in Y} \bigvee_{x \in X} \left( |A_{1f}^*(x) - A_{2f}^*(x)| + |A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y)| \right) \\
& \leq \bigvee_{y \in Y} \bigvee_{x \in X} \left( |A_{1f}^*(x) - A_{2f}^*(x)| + |A_{1-f}(x) - A_{2-f}(x)| + |B_{1-f}(y) - B_{2-f}(y)| \right) \\
& \leq \bigvee_{x \in X} |A_{1f}^*(x) - A_{2f}^*(x)| + \bigvee_{x \in X} |A_{1f}(x) - A_{2f}(x)| + \bigvee_{y \in Y} |B_{1f}(y) - B_{2f}(y)|. \quad (34)
\end{aligned}$$

Thus it follows from (13), (32) and (34) that

$$\begin{aligned}
& d_{N_2}(B_1^*, B_2^*) \\
& \leq \frac{1}{2} \left( \bigvee_{x \in X} |A_{1t}^*(x) - A_{2t}^*(x)| + \bigvee_{x \in X} |A_{1f}^*(x) - A_{2f}^*(x)| + \bigvee_{x \in X} |A_{1t}(x) - A_{2t}(x)| \right) \\
& \quad + 2 \bigvee_{x \in X} |A_{1f}(x) - A_{2f}(x)| + \bigvee_{y \in Y} |B_{1t}(y) - B_{2t}(y)| + 2 \bigvee_{y \in Y} |B_{1f}(y) - B_{2f}(y)| \\
& \leq \frac{1}{2} \left( \bigvee_{x \in X} |A_{1t}^*(x) - A_{2t}^*(x)| + \bigvee_{x \in X} |A_{1f}^*(x) - A_{2f}^*(x)| + 2 \bigvee_{x \in X} |A_{1t}(x) - A_{2t}(x)| \right) \\
& \quad + 2 \bigvee_{x \in X} |A_{1f}(x) - A_{2f}(x)| + 2 \bigvee_{y \in Y} |B_{1t}(y) - B_{2t}(y)| + 2 \bigvee_{y \in Y} |B_{1f}(y) - B_{2f}(y)| \\
& = 2d_{N_2}(A_1, A_2) + 2d_{N_2}(B_1, B_2) + d_{N_2}(A_1^*, A_2^*) \\
& \leq 2\delta_1 + 2\delta_2 + \delta_3.
\end{aligned}$$

This completes the proof.  $\square$

**Theorem 3.13.** *Let  $A_1^*$  and  $A_2^*$  be the triple implication method of IFMT ( $A_1, B_1, B_1^*$ ) and IFMT( $A_2, B_2, B_2^*$ ) based on the Lukasiewicz implication, respectively. Suppose that  $d_{N_1}(A_1, A_2) \leq \delta_1$ ,  $d_{N_1}(B_1, B_2) \leq \delta_2$ ,  $d_{N_1}(B_1^*, B_2^*) \leq \delta_3$ , then*

$$d_{N_1}(A_1^*, A_2^*) \leq 4\delta_1 \bigvee 4\delta_2 \bigvee 2\delta_3.$$

*Proof.* We firstly take the natural distance between the membership functions of  $A_1^*$  and  $A_2^*$  into consideration, by definition 2.3 and eq.(7) we have

$$\begin{aligned}
& d_n(A_{1t}^*, A_{2t}^*) \\
& = \bigvee_{x \in X} \left| \bigwedge_{y \in Y} \left\{ \left( \left( (A_{1t}(x) \rightarrow B_{1t}(y)) \bigwedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right) \rightarrow B_{1t}^*(y) \right) \right. \right. \\
& \quad \bigwedge \left. \left( (A_{1-f}(x) \rightarrow B_{1-f}(y)) \rightarrow B_{1-f}^*(y) \right) \right\} \\
& \quad - \bigwedge_{y \in Y} \left\{ \left( \left( (A_{2t}(x) \rightarrow B_{2t}(y)) \bigwedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right) \rightarrow B_{2t}^*(y) \right) \right. \\
& \quad \left. \bigwedge \left( (A_{2-f}(x) \rightarrow B_{2-f}(y)) \rightarrow B_{2-f}^*(y) \right) \right\} \right|. \quad (35)
\end{aligned}$$

Using Lemma 3.1(ii) twice, it can be easily obtained that

$$\begin{aligned}
& d_n(A_{1t}^*, A_{2t}^*) \\
& \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( \left| \left( (A_{1t}(x) \rightarrow B_{1t}(y)) \wedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right) \rightarrow B_{1t}^*(y) \right. \right. \\
& \quad \left. \left. - \left( (A_{2t}(x) \rightarrow B_{2t}(y)) \wedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right) \rightarrow B_{2t}^*(y) \right| \right. \\
& \quad \left. \bigvee \left| (A_{1-f}(x) \rightarrow B_{1-f}(y)) \rightarrow B_{1-f}^*(y) \right. \right. \\
& \quad \left. \left. - (A_{2-f}(x) \rightarrow B_{2-f}(y)) \rightarrow B_{2-f}^*(y) \right| \right). \tag{36}
\end{aligned}$$

Then it follows Lemma 3.3(ii) that

$$\begin{aligned}
& d_n(A_{1t}^*, A_{2t}^*) \\
& \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( 2 \left| (A_{1t}(x) \rightarrow B_{1t}(y)) \wedge (A_{1-f}(x) \rightarrow B_{1-f}(y)) \right. \right. \\
& \quad \left. \left. - (A_{2t}(x) \rightarrow B_{2t}(y)) \wedge (A_{2-f}(x) \rightarrow B_{2-f}(y)) \right| \bigvee 2 \left| B_{1t}^*(y) - B_{2t}^*(y) \right| \right. \\
& \quad \left. \bigvee 2 \left| A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y) \right| \right. \\
& \quad \left. \bigvee 2 \left| B_{1-f}^*(y) - B_{2-f}^*(y) \right| \right). \tag{37}
\end{aligned}$$

According to Lemma 3.1(ii), we further have

$$\begin{aligned}
& d_n(A_{1t}^*, A_{2t}^*) \\
& \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( 2 \left| B_{1t}^*(y) - B_{2t}^*(y) \right| \bigvee 2 \left| A_{1t}(x) \rightarrow B_{1t}(y) - A_{2t}(x) \rightarrow B_{2t}(y) \right| \right. \\
& \quad \left. \bigvee 2 \left| A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y) \right| \bigvee 2 \left| B_{1-f}^*(y) - B_{2-f}^*(y) \right| \right. \\
& \quad \left. \bigvee 2 \left| A_{1-f}(x) \rightarrow B_{1-f}(y) - A_{2-f}(x) \rightarrow B_{2-f}(y) \right| \right). \tag{38}
\end{aligned}$$

For (38), using Lemma 3.3(ii) again, we can obtain

$$\begin{aligned}
& d_n(A_{1t}^*, A_{2t}^*) \\
& \leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( 2 \left| B_{1t}^*(y) - B_{2t}^*(y) \right| \bigvee 4 \left| A_{1t}(x) - A_{2t}(x) \right| \bigvee 4 \left| B_{1t}(y) - B_{2t}(y) \right| \right. \\
& \quad \left. \bigvee 4 \left| A_{1-f}(x) - A_{2-f}(x) \right| \bigvee 4 \left| B_{1-f}(y) - B_{2-f}(y) \right| \bigvee 2 \left| B_{1-f}^*(y) - B_{2-f}^*(y) \right| \right) \\
& = \bigvee_{x \in X} \bigvee_{y \in Y} \left( 4 \left| A_{1t}(x) - A_{2t}(x) \right| \bigvee 4 \left| A_{1-f}(x) - A_{2-f}(x) \right| \bigvee 4 \left| B_{1t}(y) - B_{2t}(y) \right| \right. \\
& \quad \left. \bigvee 4 \left| B_{1-f}(y) - B_{2-f}(y) \right| \bigvee 2 \left| B_{1t}^*(y) - B_{2t}^*(y) \right| \bigvee 2 \left| B_{1-f}^*(y) - B_{2-f}^*(y) \right| \right). \tag{39}
\end{aligned}$$

Secondly, we consider the natural distance between the non-membership functions of  $A_1^*$  and  $A_2^*$ , by Definition 2.3 and eq.(7) we have

$$d_n(A_{1f}^*, A_{2f}^*) = \bigvee_{x \in X} \left| \bigvee_{y \in Y} \{1 - (A_{1-f}(x) \rightarrow B_{1-f}(y)) \rightarrow B_{1-f}^*(y)\} - \bigvee_{y \in Y} \{1 - (A_{2-f}(x) \rightarrow B_{2-f}(y)) \rightarrow B_{2-f}^*(y)\} \right|. \quad (40)$$

Then using the similar method above, it can be easily gotten from (40) that

$$d_n(A_{1f}^*, A_{2f}^*) \leq \bigvee_{x \in X} \bigvee_{y \in Y} (4|A_{1f}(x) - A_{2f}(x)| \bigvee 4|B_{1f}(y) - B_{2f}(y)| \bigvee 2|B_{1f}^*(y) - B_{2f}^*(y)|). \quad (41)$$

Thus it follows from (8), (39) and (41) that

$$\begin{aligned} d_{N_1}(A_1^*, A_2^*) &= d_n(A_{1t}^*, A_{2t}^*) \bigvee d_n(A_{1f}^*, A_{2f}^*) \\ &\leq \bigvee_{x \in X} \bigvee_{y \in Y} \left( 4|A_{1t}(x) - A_{2t}(x)| \bigvee 4|A_{1f}(x) - A_{2f}(x)| \bigvee 4|B_{1t}(y) - B_{2t}(y)| \right. \\ &\quad \left. \bigvee 4|B_{1f}(y) - B_{2f}(y)| \bigvee 2|B_{1t}^*(y) - B_{2t}^*(y)| \bigvee 2|B_{1f}^*(y) - B_{2f}^*(y)| \right). \\ &= 4d_{N_1}(A_1, A_2) \bigvee 4d_{N_1}(B_1, B_2) \bigvee 2d_{N_1}(B_1^*, B_2^*) \\ &\leq 4\delta_1 \bigvee 4\delta_2 \bigvee 2\delta_3. \end{aligned}$$

This completes the proof.  $\square$

**Theorem 3.14.** *Let  $A_1^*$  and  $A_2^*$  be the triple implication solutions of IFMT  $(A_1, B_1, B_1^*)$  and IFMT  $(A_2, B_2, B_2^*)$  based on the Lukasiewicz implication, respectively. Suppose that  $d_{N_2}(A_1, A_2) \leq \delta_1$ ,  $d_{N_2}(B_1, B_2) \leq \delta_2$ ,  $d_{N_2}(B_1^*, B_2^*) \leq \delta_3$ , then*

$$d_{N_2}(A_1^*, A_2^*) \leq 2\delta_1 + 2\delta_2 + \delta_3.$$

*Proof.* The proof is similar to that of above Theorems.  $\square$

**Remark 3.15.** Suppose that  $\delta_i$  is close to 0 ( $i=1,2,3$ ), then it follows from Theorem 3.11 and Theorem 3.12 that both  $d_{N_1}(B_1^*, B_2^*)$  and  $d_{N_2}(B_1^*, B_2^*)$  are close to 0. Based on this fact, we can see that small disturbances on the inputs  $A, B$  and  $A^*$  imply small changes of the output  $B^*$  of IFMP( $A, B, A^*$ ), hence the triple implication method of Intuitionistic Fuzzy Modus Ponens has a good behavior of robustness under two kinds of natural distances given by the present paper. The same conclusion can be drawn with respect to the triple implication method of Intuitionistic Fuzzy Modus Tollens.

**Remark 3.16.** It is obvious that if  $g$  is reductive, then  $B$  is the solution of IFMP  $(A, B, A)$ , therefore the continuity (i.e., approximation property) of  $g$  is just right its robustness in the case that  $A_1 = A_2 = A, B_1 = B_2 = B$  and  $A_1^* = A_2^* = A$ . In this case, both the continuity (i.e., approximation property) and the robustness reflect that small disturbances on the input do not lead to large deviations of the output.

A similar analysis can be done about the method  $h$  for Intuitionistic Fuzzy Modus Tollens.

**Remark 3.17.** It is worth noting that both the present paper and references [24, 38] above have discussed the robustness of fuzzy reasoning, but the robustness of fuzzy connectives and fuzzy reasoning in the reference [24] is discussed through the sensitivity of interval-valued fuzzy connectives, the robustness of intuitionistic fuzzy connectives in fuzzy reasoning is discussed in the reference [38] through the evaluation of sensitivity in  $n$ -order function on the class of intuitionistic fuzzy sets, while the present is through some distances defined on the set of all intuitionistic fuzzy sets to discuss the continuity, approximation property and robustness of the triple implication method of intuitionistic fuzzy reasoning.

#### 4. Conclusions

In this paper, we discussed the continuity, approximation property and robustness of triple implication method for Intuitionistic Fuzzy Modus Ponens model by using two kinds of natural distances based on Łukasiewicz implication, and for intuitionistic fuzzy modus tollens model, we use similar method to take continuity and robustness into consideration. Our results showed that both the triple implication methods for Intuitionistic Fuzzy Modus Ponens and Intuitionistic Fuzzy Modus Tollens have continuity and good behavior of robustness. Of course, we should note that the robustness results presented in this paper are conservative in certain sense. A fuzzy knowledge base normally contains many fuzzy rules, as in the case of fuzzy control. We can treat inference with multiple rules as a generalized form of generalized modus ponens. However, since multiple rules are involved, we may obtain different robustness results between combination based inference and individual-rule based inference. This is a problem left for further investigation. Whether triple implication method for intuitionistic fuzzy modus tollens model is of approximation property is another problem left. Finally, the present paper is mainly based on intuitionistic fuzzy implication operator induced by Łukasiewicz implication operator, the results about other intuitionistic fuzzy implication operators are possible topics for future study.

**Acknowledgements.** Authors would like to express their sincere thanks to the valuable comments and suggestions from editors and reviewers. This work was supported by the National Natural Science Foundation of China (No. 11261032).

#### REFERENCES

- [1] K. Atanassov, *Geometrical interpretation of the elements of the intuitionistic fuzzy objects*, Preprint IM-MFAIS-1-89, Sofia, 1989. Reprinted: International Journal of Bioautomation, **20(1)**(2016), S43-S54.
- [2] K. Atanassov, *Intuitionistic fuzzy sets*, Fuzzy Sets and Systems, **20(1)**(1986), 87-96.
- [3] K. Atanassov, *Intuitionistic fuzzy sets*, International Journal of Bioautomation, **20(S1)**(2016), S27-S42.
- [4] K. Atanassov, *Intuitionistic fuzzy sets*. Springer, Heidelberg, 1999.
- [5] K. Atanassov, *On intuitionistic fuzzy sets theory*, Springer, Berlin, **8(1)**(2012), 147-163.

- [6] I. Bloch, *Mathematical morphology on bipolar fuzzy sets: General algebraic framework*, International Journal of Approximate Reasoning, **53(7)**(2012), 1031-1060.
- [7] F. Chen, W. H. Xu, C. Z. Bai and X. Gao, *A novel approach to guarantee good robustness of fuzzy reasoning*, Applied Soft Computing, **41(C)**(2016), 224-234.
- [8] R. Cignoli, I. M. L. Dottaviano and D. Mundici, *Algebraic Foundations of Many-Valued Reasoning*, Kluwer Academic Publishers, Dordrecht, 2000.
- [9] C. Cornelis, G. Deschrijver and E. E. Kerre, *Implication in intuitionistic fuzzy and interval-valued fuzzy set theory: construction, classification, application*, International Journal of Approximate Reasoning, **35(1)**(2004), 55-95.
- [10] S. S. Dai, D. W. Pei and D. Guo, *Robustness analysis of full implication inference method*, International Journal of Approximate Reasoning, **54(5)**(2013), 653-666.
- [11] S. S. Dai, D. W. Pei and S. M. Wang, *Perturbation of fuzzy sets and fuzzy reasoning based on normalized Minkowski distances*, Fuzzy Sets and Systems, **189(1)**(2012), 63-73.
- [12] S. Das, B. Dutta and D. Guha, *Weight computation of criteria in a decision-making problem by knowledge measure with intuitionistic fuzzy set and interval-valued intuitionistic fuzzy set*, Soft Computing, **20(9)**(2016), 3421-3442.
- [13] G. Deschrijver, C. Cornelis and E. E. Kerre, *Class of intuitionistic fuzzy t-norms satisfying the residuation principle*, International Journal of Uncertainty Fuzziness and Knowledge-Based Systems, **11(6)**(2003), 691-709.
- [14] G. Deschrijver, C. Cornelis and E. E. Kerre, *On the representation of intuitionistic fuzzy t-norms and t-conorms*, IEEE Transactions on Fuzzy Systems, **12(1)**(2004), 45-61.
- [15] J. Y. Duan and Y. M. Li, *Robustness analysis of logic metrics on  $F(X)$* , International Journal of Approximate Reasoning, **61**(2015), 33-42.
- [16] D. Dubois and H. Prade, *Fuzzy Sets in approximate reasoning*, Fuzzy Sets and Systems, **40(1)**(1991), 143-244.
- [17] D. Dubois and H. Prade, *Gradualness, uncertainty and bipolarity: Making sense of fuzzy sets*, Fuzzy Sets and Systems, **192(3)**(2012), 3-24.
- [18] F. Esteva and L. Godo, *Monoidal t-norm based logic: Towards a logic for left-continuous t-norms*, Fuzzy Sets and Systems, **124(3)**(2001), 271-288.
- [19] C. Franco, J. Montero and J. T. Rodriguez, *A fuzzy and bipolar approach to preference modeling with application to need and desire*, Fuzzy Sets and Systems, **214(1)**(2013), 20-34.
- [20] P. Hajek, *Mathematics of Fuzzy Logic*, Kluwer Academic Publishers, Dordrecht, 1998.
- [21] X. X. He, K. Y. Qin and Y. F. Li, *Robustness of fuzzy connectives and fuzzy reasoning*, Fuzzy Sets and Systems, **225(12)**(2013), 93-105.
- [22] D. H. Hong and S. Y. Hwang, *A note on the value similarity of fuzzy systems variables*, Fuzzy Sets and Systems, **66(3)**(1994), 383-386.
- [23] C. C. Lee, *Fuzzy logic in control systems: Fuzzy logic controller*, IEEE Transactions on Systems, Man, and Cybernetics, **20(2)**(1990), 404-418.
- [24] D. C. Li, Y. M. Li and Y. J. Xie, *Robustness of interval-valued fuzzy inference*, Information Sciences, **181(20)**(2011), 4754-4764.
- [25] D. F. Li and C. T. Cheng, *New similarity measures of intuitionistic fuzzy sets and application to pattern recognitions*, Pattern Recognition Letters, **23(1-3)**(2002), 221-225.
- [26] Y. F. Li, K. Y. Qin, X. X. He and D. Meng, *Robustness of fuzzy connectives and fuzzy reasoning with respect to general divergence measures*, Fuzzy Sets and Systems, **294**(2016), 63-78.
- [27] H. W. Liu and G. J. Wang, *Continuity of triple I methods based on several implications*, Mathematical and Computer Modelling, **56(8)**(2008), 2079-2087.
- [28] H. W. Liu and G. J. Wang, *Multi-criteria decision-making methods based on intuitionistic fuzzy sets*, European Journal of Operational Research, **179(1)**(2007), 220-233.
- [29] H. W. Liu and G. J. Wang, *Unified forms of fully implicational restriction methods for fuzzy reasoning*, Information Sciences, **177(3)**(2007), 956-966.
- [30] H. W. Liu, *New similarity measures between intuitionistic fuzzy sets and between elements*, Mathematical and Computer Modelling, **42(1)**(2005), 61-70.

- [31] R. Lowen and W. Peeters, *Distances between fuzzy sets representing grey level images*, Fuzzy Sets and Systems, **99(2)**(1998) 135-149.
- [32] E. H. Mamdani and B. R. Gaines, *Fuzzy Reasoning and its Applications*, Academic Press, London, **18(11)**(1981) 1925-1935.
- [33] D. W. Pei, *Formalization of implication based fuzzy reasoning method*, International Journal of Approximate Reasoning, **53(5)**(2012), 837-846.
- [34] D. W. Pei, *On equivalent forms of fuzzy logic systems NM and IMTL*, Fuzzy Sets and Systems, **138(1)**(2003), 187-195.
- [35] D. W. Pei, *On the strict logic foundation of fuzzy reasoning*, Soft Computing, **8(8)**(2004), 539-545.
- [36] D. W. Pei, *The full implication triple I algorithms and their consistency in fuzzy reasoning*, Journal of Mathematical Research and Exposition, **24(2)**(2004), 359-368 (in Chinese).
- [37] D. W. Pei, *Unified full implication algorithms of fuzzy reasoning*, Information Sciences, **178(2)**(2008), 520-530.
- [38] R. H. S. Reiser and B. Bedregal, *Robustness on intuitionistic fuzzy connectives*, Tema, **15(2)**(2014), 133-149.
- [39] R. Srinivasan and S. S. Begum, *Properties on Intuitionistic Fuzzy Sets of Third Type*, Fuzzy Information Processing Society, IEEE, **12(2)**(2017), 189-195.
- [40] Y. M. Tang and X. Z. Yang, *Symmetric implicational method of fuzzy reasoning*, International Journal of Approximate Reasoning, **54(8)**(2015), 1034-1048.
- [41] G. J. Wang and J. Y. Duan, *On robustness of the full implication triple I inference method with respect to finer measurements*, International Journal of Approximate Reasoning, **55(3)**(2014), 787-796.
- [42] G. J. Wang and L. Fu, *Unified forms of triple I method*, Computers and Mathematics with Applications, **49(5-6)**(2005), 923-932.
- [43] G. J. Wang and H. Wang, *Non-fuzzy versions of fuzzy reasoning in classical logics*, Information Sciences, **138(1)**(2001), 211-236.
- [44] G. J. Wang, *Formalized theory of general fuzzy reasoning*, Information Sciences, **160(1)**(2004), 251-266.
- [45] G. J. Wang, *Full implicational triple I method for fuzzy reasoning*, Science in China, Series E, **29(1)**(1999), 43-53.
- [46] G. J. Wang, *Introduction to Mathematical Logic and Resolution Principle*, Science Press, Beijing, 2006 (in Chinese).
- [47] G. J. Wang, *Non-classical Mathematical Logic and Approximate Reasoning*, Science Press, Beijing, 2008 (in Chinese).
- [48] G. J. Wang, *On the logic foundation of fuzzy reasoning*, Information Sciences, **117(1-2)**(1999), 47-88.
- [49] L. X. Wang, *A Course in fuzzy systems and control*, Upper Saddle River, Prentice Hall PTR, NJ, 1997.
- [50] W. H. Xu, Z. K. Xie, J. Y. Yang and Y. E. You-Pei, *Continuity and approximation properties of two classes of algorithms for fuzzy inference*, Journal of Software, **15(10)**(2004), 1485-1492 (in chinese).
- [51] L. A. Zadeh, *Fuzzy sets, information and control*, Information and Control, **8(3)**(1965), 338-353.
- [52] L. A. Zadeh, *Generalized theory of uncertainty (GTU)-principal concepts and ideas*, Computational Statistics and Data Analysis, **51(1)**(2006), 15-46.
- [53] L. A. Zadeh, *Outline of a new approach to the analysis of complex systems and decision processes*, IEEE Transactions on Systems, Man, and Cybernetics, **smc-3(1)**(1973), 28-44.
- [54] L. A. Zadeh, *Position Paper: Toward extended fuzzy logic-A first step*, Fuzzy Sets and Systems, **160(21)**(2009), 3175-3181.
- [55] L. A. Zadeh, *The concept of a linguistic variable and its application to approximate reasoning*, Learning Systems and Intelligent Robots. Springer US, **8(3)**(1974), 199-249.

- [56] M. C. Zheng, Z. K. Shi and Y. Liu, *Triple I method of approximate reasoning on Atanassov's intuitionistic fuzzy sets*, International Journal of Approximate Reasoning, **55(6)**(2014), 1369-1382.
- [57] M. C. Zheng, Z. K. Shi and Y. Liu, *Triple I method of intuitionistic fuzzy reasoning based on residual implicator*, Information Sciences, **43(6)**(2013), 810-820 (in chinese).

JUN LI\*, SCHOOL OF SCIENCE, LANZHOU UNIVERSITY OF TECHNOLOGY, LANZHOU 730050,  
GANSU, CHINA  
*E-mail address:* lijun@lut.cn

YAN LIU, SCHOOL OF SCIENCE, LANZHOU UNIVERSITY OF TECHNOLOGY, LANZHOU 730050,  
GANSU, CHINA  
*E-mail address:* 774558749@qq.com

\*CORRESPONDING AUTHOR