

Stability problem for Pexiderized Cauchy-Jensen type functional equations of fuzzy number-valued mappings

M. Miah¹, F. Mirzaee² and H. Khodaei³

^{1,2,3}Faculty of Mathematical Sciences and Statistics, Malayer University, P.O. Box 65719-95863, Malayer, Iran

minamiahi@gmail.com, f.mirzaee@malayeru.ac.ir, hkhodaei@malayeru.ac.ir

Abstract

We investigate the stability problems of the n -dimensional Cauchy-Jensen type and the n -dimensional Pexiderized Cauchy-Jensen type fuzzy number-valued functional equations in Banach spaces by using the metric defined on a fuzzy number space. Under some suitable conditions, some properties of the solutions for these equations such as existence and uniqueness are discussed. Our results can be regarded as important extensions of stability results corresponding to single-valued functional equations and set-valued functional equations, respectively.

Keywords: Fuzzy number-valued mapping, stability, Pexiderized Cauchy-Jensen equation, fuzzy analysis.

1 Introduction

In set-valued analysis, a functional inclusion is called stable if any function which satisfies this inclusion approximately is near to a true solution of the functional inclusion (see [7, 8, 15, 21, 37]). Nikodem and Popa [28] considered the general solution of set-valued functions satisfying linear inclusion relation, which can be regarded as a generalization of the additive single-valued functional equation. By means of the inclusion relation, Park et al. [23, 30] investigated the stability of some set-valued functional equations. Jang et al. [19] and Chu et al. [13] studied the stability of an n -dimensional additive set-valued functional equation and an n -dimensional cubic set-valued functional equation, respectively.

Some interesting results concerning the Cauchy-Jensen functional equation

$$f\left(\frac{x+y}{2}\right) + f\left(\frac{x-y}{2}\right) = f(x),$$

and the general linear functional equation with real constants a, b, A, B ,

$$f(ax + by) = Af(x) + Bf(y), \tag{1}$$

have been obtained in [6, 17, 20, 23, 24, 29, 32, 39]. In particular, the equation (1) includes the linear equation for $A = a$ and $B = b$. If $t \in (0, 1)$, $a = A = t$ and $b = B = 1 - t$, then (1) has the form

$$f(tx + (1-t)y) = tf(x) + (1-t)f(y),$$

and its solution is called an affine function (see, among others, [29, 33]). Clearly, the equation (1) also includes the Cauchy equation for $a = b = A = B = 1$ and the Jensen equation for $a = b = A = B = \frac{1}{2}$.

The functional equation

$$f(x + y) = g(x) + h(y),$$

with three unknown functions f, g, h treated by Pexider [31], which is known as the Pexiderized Cauchy equation. In the case $f = g = h$, this equation reduces to Cauchy equation. The Pexiderized Cauchy equation has a simple interpretation: The value of f at $x + y$ is the sum of two values, of which one depends only on x , and the other only on y (separation of variables).

In 1991, Nikodem [27] investigated the stability of the Pexider Cauchy equation. This paper seems to be the first one concerning the stability problem of the Pexider Cauchy equation. Solutions and the stability of various Pexiderized Cauchy and Pexiderized Cauchy-Jensen functional equations in several variables have been investigated by several authors (see, among others, [12, 14, 16, 22, 25, 26]).

The stability of the Cauchy-Jensen and the general linear type fuzzy number-valued functional equations in Banach spaces by using the metric defined on a fuzzy number space investigated by Wu and Jin [39], which are generalizations of the main results obtained in [17, 23]. We refer the reader to [1, 5, 9, 10, 11, 35, 36, 38] for some recent works and discussions on this topic.

The purpose of this paper is to extend the single-valued and set-valued functional equations to fuzzy number-valued functional equations and investigate the stability problems of the n -dimensional Cauchy-Jensen type fuzzy number-valued functional equation

$$f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) = f(x_1). \quad (2)$$

and the n -dimensional Pexiderized Cauchy-Jensen type fuzzy number-valued functional equation

$$f_1\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{j=2}^n f_j\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right) = f_{n+1}(x_1), \quad (3)$$

in Banach spaces via the metric defined on a fuzzy number space. Under some suitable conditions, some properties of the solutions for these equations such as existence and uniqueness are discussed. Notice that the supremum metric, as a generalization of the Hausdorff metric, is applied to characterize the fuzzy number-valued functional inequality. It has been shown that the stability of Cauchy, Jensen, Cauchy-Jensen, Pexiderized Cauchy, Pexiderized Jensen, Pexiderized affine, Pexiderized linear and Pexiderized general linear type fuzzy number-valued functional equations can be obtained as special cases from our results. Therefore, the results obtained in the present paper improve and extend the corresponding results in previous works [12, 17, 23, 27, 30, 39].

2 Preliminaries

In this section, we consider some basic concepts and results (see [3, 39]) which will be used in the sequel. From now on, assume that X, Y are Banach spaces and B is a subspace of Y .

Suppose a function $u : X \rightarrow [0, 1]$ satisfies the following conditions:

1. $[u]^\alpha = \{x \in X : u(x) \geq \alpha\}$ is a non-empty compact convex subset of X , for all $\alpha \in (0, 1]$,
2. $[u]^0 = \overline{\{x : u(x) > 0\}}$ is compact, where \overline{A} denotes the closure of A .

Then u is called a fuzzy number on X . The set of all fuzzy numbers on X is denoted by X_F .

For $u, v \in X_F$ and $\lambda \in \mathbb{R}$, by Zadeh extension principle, we obtain the following properties about addition $u + v$ and scalar multiplication $\lambda \cdot u$ (see [3, 39]):

- (i) $[u + v]^\alpha = [u]^\alpha + [v]^\alpha$,
- (ii) $[\lambda \cdot u]^\alpha = \lambda[u]^\alpha$.

Note that $\tilde{0}$ is the zero element in X_F ; i.e., $\tilde{0} + u = u + \tilde{0} = u$, for all $u \in X_F$.

Defining the mapping $D : X_F \times X_F \rightarrow [0, \infty)$ by

$$D(u, v) = \sup_{\alpha \in [0, 1]} d_H([u]^\alpha, [v]^\alpha),$$

where d_H is Hausdorff metric. Then D satisfies the following properties:

- (P1) $D(\lambda \cdot u, \lambda \cdot v) = |\lambda|D(u, v)$,
- (P2) $D(u + w, v + w) = D(u, v)$,
- (P3) $D(u + v, w + e) \leq D(u, w) + D(v, e)$

for all $\lambda \in \mathbb{R}$ and $u, v, w, e \in X_F$, and (X_F, D) is a complete metric space.

Let $u, v \in X_F$. If there exists $w \in X_F$ such that $u = v + w$, then w is called the Hukuhara difference (H-difference) of u and v , and it is denoted by $u - v$ (see [18, 34]).

Notation 2.1. Let $\lambda \in \mathbb{R}$ and $u, v \in \mathbb{R}_F$. If we denote $\|u\|_F = D(u, \tilde{0})$, then $\|\cdot\|_F$ has the properties of a usual norm on \mathbb{R}_F ; i.e., $\|u\|_F = 0$ iff $u = \tilde{0}$, $\|\lambda \cdot u\|_F = |\lambda| \|u\|_F$ and $\|u + v\|_F \leq \|u\|_F + \|v\|_F$. Moreover, note that $(\mathbb{R}_F, +, \cdot)$ is not a linear space over \mathbb{R} and consequently $(\mathbb{R}_F, \|\cdot\|_F)$ cannot be a normed space (see [2] and [4, Theorem 1 and Remark 2]).

3 Stability of the Cauchy-Jensen type fuzzy number-valued functional equation in n variables

In this section, the stability of the equation (2) is established, in which f indicates a fuzzy number-valued mapping. Several special cases are also considered.

Theorem 3.1. Let $f : B \rightarrow X_F$ be a fuzzy number-valued mapping such that

$$D\left(f\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right), f(x_1)\right) < \varepsilon, \quad (4)$$

for some $\varepsilon > 0$ and for all $x_1, \dots, x_n \in B$. Then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that

$$D(f(x), \mathcal{G}(x)) \leq \frac{\varepsilon}{n-1},$$

for all $x \in B$.

Proof. Setting $x_1 = nx$ and $x_i = 0$ ($2 \leq i \leq n$) in (4), we obtain

$$D\left(f(x), \frac{1}{n}f(nx)\right) < \frac{\varepsilon}{n},$$

for all $x \in B$. Then, by induction on m , we have

$$D\left(f(x), \frac{1}{n^m}f(n^m x)\right) < \sum_{j=1}^m \left(\frac{1}{n}\right)^j \varepsilon. \quad (5)$$

Let $f_0(x) = f(x)$ and $f_m(x) = \frac{1}{n^m}f(n^m x)$, where $m \in \mathbb{N}$ and $x \in B$. Then, by (5) and the properties (P2) and (P3) of D , we obtain

$$\begin{aligned} D(f_m(x), f_{m-1}(x)) &= D\left(\frac{1}{n^m}f(n^m x) + f(x), f(x) + \frac{1}{n^{m-1}}f(n^{m-1}x)\right) \\ &\leq D\left(\frac{1}{n^m}f(n^m x), f(x)\right) + D\left(f(x), \frac{1}{n^{m-1}}f(n^{m-1}x)\right) \\ &< \sum_{j=1}^m \left(\frac{1}{n}\right)^j \varepsilon + \sum_{j=1}^{m-1} \left(\frac{1}{n}\right)^j \varepsilon, \end{aligned}$$

for all $x \in B$. Thus the sequence $\{f_m(x)\}$ is Cauchy in X_F . Since (X_F, D) is complete, we can define $\mathcal{G}(x) = \lim_{m \rightarrow \infty} f_m(x)$ for any $x \in B$.

The next step is to show that \mathcal{G} is additive. By (4), we can conclude that

$$\begin{aligned} &D\left(f_m\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{j=2}^n f_m\left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n}\right), f_m(x_1)\right) \\ &= D\left(\frac{1}{n^m}f\left(\frac{\sum_{i=1}^n n^m x_i}{n}\right) + \frac{1}{n^m} \sum_{j=2}^n f\left(\frac{\sum_{i=1, i \neq j}^n n^m x_i - n^m(n-1)x_j}{n}\right), \frac{1}{n^m}f(n^m x_1)\right) < \frac{\varepsilon}{n^m}, \end{aligned}$$

for all $x_1, \dots, x_n \in B$. Hence,

$$\lim_{m \rightarrow \infty} D \left(f_m \left(\frac{\sum_{i=1}^n x_i}{n} \right) + \sum_{j=2}^n f_m \left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n} \right), f_m(x_1) \right) = 0.$$

By the continuity of the metric D , we obtain

$$D \left(\mathcal{G} \left(\frac{\sum_{i=1}^n x_i}{n} \right) + \sum_{j=2}^n \mathcal{G} \left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n} \right), \mathcal{G}(x_1) \right) = 0,$$

meaning

$$\mathcal{G} \left(\frac{\sum_{i=1}^n x_i}{n} \right) + \sum_{j=2}^n \mathcal{G} \left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n} \right) = \mathcal{G}(x_1), \quad (6)$$

for all $x_1, \dots, x_n \in B$. Letting $x_i = 0$ ($1 \leq i \leq n$) in (6), we obtain $\mathcal{G}(0) = \tilde{0}$. Setting $x_1 = x$ and $x_i = 0$ ($2 \leq i \leq n$) in (6), we get

$$n\mathcal{G} \left(\frac{x}{n} \right) = \mathcal{G}(x), \quad (7)$$

for all $x \in B$. Setting $x_i = 0$ ($3 \leq i \leq n$) in (6) and using (7), we have

$$\frac{n-1}{n}\mathcal{G}(x_1 + x_2) + \frac{1}{n}\mathcal{G}(x_1 - (n-1)x_2) = \mathcal{G}(x_1), \quad (8)$$

for all $x_1, x_2 \in B$. Putting $x_1 = x_1 + (n-1)x_2$ in (8), one finds

$$\frac{n-1}{n}\mathcal{G}(x_1 + nx_2) + \frac{1}{n}\mathcal{G}(x_1) = \mathcal{G}(x_1 + (n-1)x_2), \quad (9)$$

for all $x_1, x_2 \in B$. Replacing x_1 by 0 and x_2 by x into (9) and using (7), one gets

$$\mathcal{G}((n-1)x) = (n-1)\mathcal{G}(x), \quad (10)$$

for all $x \in B$. Replacing x_1 by 0 and x_2 by x into (8) and using (10), we obtain $\mathcal{G}(-x) = -\mathcal{G}(x)$ for all $x \in B$; that is, \mathcal{G} is odd. Setting $x_2 = x_2 - x_1$ in (8), we get

$$\frac{n-1}{n}\mathcal{G}(x_2) + \frac{1}{n}\mathcal{G}(nx_1 - (n-1)x_2) = \mathcal{G}(x_1), \quad (11)$$

for all $x_1, x_2 \in B$. Replacing x_1 by $\frac{x_1}{n}$ and x_2 by $-\frac{x_2}{n-1}$ into (11) and using (7), (10) and the oddness of \mathcal{G} , we have

$$\mathcal{G}(x_1 + x_2) = \mathcal{G}(x_1) + \mathcal{G}(x_2),$$

for all $x_1, x_2 \in B$; that is, \mathcal{G} is additive.

By letting $m \rightarrow \infty$ in (5), we immediately achieve $D(f(x), \mathcal{G}(x)) \leq \frac{\varepsilon}{n-1}$ for all $x \in B$.

Finally, we prove the uniqueness of \mathcal{G} . Assume that there exist additive mappings $\mathcal{G}_1, \mathcal{G}_2 : B \rightarrow X_F$ satisfying the last inequality. Then,

$$\begin{aligned} D(\mathcal{G}_1(x), \mathcal{G}_2(x)) &= \frac{1}{m} D(m\mathcal{G}_1(x), m\mathcal{G}_2(x)) \\ &\leq \frac{1}{m} (D(\mathcal{G}_1(mx), f(mx)) + D(f(mx), n\mathcal{G}_2(mx))) \\ &< \frac{2\varepsilon}{m(n-1)}, \end{aligned}$$

for all $x \in B$ and for any $m \in \mathbb{N}$. Since the right-hand side of the last inequality tends to zero as $m \rightarrow \infty$, we conclude that $\mathcal{G}_1(x) = \mathcal{G}_2(x)$ for all $x \in B$, as desired. \square

From Theorem 3.1, we easily obtain the following.

Corollary 3.2. ([39, Theorem 1]) *If a fuzzy number-valued mapping $f : B \rightarrow X_F$ satisfies the inequality (4) with $n = 2$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that*

$$D(f(x), \mathcal{G}(x)) \leq \varepsilon, \quad (12)$$

for all $x \in B$.

In the next result, we establish the stability of the Cauchy type fuzzy number-valued functional equation.

Corollary 3.3. *Let $f : B \rightarrow X_F$ be a fuzzy number-valued mapping such that*

$$D(f(x) + f(y), f(x + y)) < \varepsilon, \quad (13)$$

for some $\varepsilon > 0$ and for all $x, y \in B$. Then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that (12) holds for all $x \in B$.

Proof. Substituting $x = \frac{x+y}{2}$ and $y = \frac{x-y}{2}$ in (13) gives

$$D\left(f\left(\frac{x+y}{2}\right) + f\left(\frac{x-y}{2}\right), f(x)\right) < \varepsilon,$$

for all $x, y \in B$. Then f satisfies the inequality (4) with $n = 2$. Then, by Theorem 3.1, we find a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that (12) holds for all $x \in B$. \square

Now, we obtain the stability of the Jensen type fuzzy number-valued functional equation.

Corollary 3.4. *If a fuzzy number-valued mapping $f : B \rightarrow X_F$ satisfies the inequality*

$$D\left(2f\left(\frac{x+y}{2}\right), f(x) + f(y)\right) < \varepsilon, \quad (14)$$

for some $\varepsilon > 0$ and for all $x, y \in B$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that

$$D(f(x) - f(0), \mathcal{G}(x)) \leq 2\varepsilon,$$

for all $x \in B$.

Proof. Let $g(x) := f(x) - f(0)$ for all $x \in B$. Then $g(0) = \tilde{0}$ and (14) yields that

$$D\left(2g\left(\frac{x+y}{2}\right), g(x) + g(y)\right) < \varepsilon, \quad (15)$$

for all $x, y \in B$. Replacing x by $x + y$ and y by 0 in (15), we get

$$D\left(2g\left(\frac{x+y}{2}\right), g(x+y)\right) < \varepsilon, \quad (16)$$

for all $x, y \in B$. It follows from (15) and (16) that

$$\begin{aligned} D(g(x+y), g(x) + g(y)) &= D\left(g(x+y) + 2g\left(\frac{x+y}{2}\right), 2g\left(\frac{x+y}{2}\right) + g(x) + g(y)\right) \\ &\leq D\left(g(x+y), 2g\left(\frac{x+y}{2}\right)\right) + D\left(2g\left(\frac{x+y}{2}\right), g(x) + g(y)\right) \\ &< 2\varepsilon, \end{aligned}$$

for all $x, y \in B$. Using the same method as in the proof of Corollary 3.3, we find a unique additive mapping $\mathcal{G} : X \rightarrow Y$ such that

$$D(g(x), \mathcal{G}(x)) \leq 2\varepsilon,$$

holds for all $x \in B$. This completes the proof of this result. \square

4 Stability of the Pexiderized Cauchy-Jensen type fuzzy number-valued functional equation in n variables

In this section, the stability of the equation (3) is established, in which f indicates a fuzzy number-valued mapping. Some special cases involving the stability of Pexiderized Cauchy, Pexiderized Jensen, Pexiderized affine, Pexiderized linear and Pexiderized general linear type fuzzy number-valued functional equations are also presented.

Theorem 4.1. *Let $f_1, \dots, f_{n+1} : B \rightarrow X_F$ be fuzzy number-valued mappings with $f_1(0) = \dots = f_n(0) = \tilde{0}$ such that*

$$D \left(f_1 \left(\frac{\sum_{i=1}^n x_i}{n} \right) + \sum_{j=2}^n f_j \left(\frac{\sum_{i=1, i \neq j}^n x_i - (n-1)x_j}{n} \right), f_{n+1}(x_1) \right) < \varepsilon, \quad (17)$$

for some $\varepsilon > 0$ and for all $x_1, \dots, x_n \in B$. Then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that

$$D(f_k(x), \mathcal{G}(x)) \leq \frac{2n}{n-1}\varepsilon, \quad (k = 1, 2, \dots, n), \quad (18)$$

$$D(f_{n+1}(x), \mathcal{G}(x)) \leq \frac{n+1}{n-1}\varepsilon, \quad (19)$$

for all $x \in B$.

Proof. Letting $x_1 = x$ and $x_i = \frac{(n-2)x+ny}{2n-2}$ ($2 \leq i \leq n$) in (17), we get

$$D \left(f_1 \left(\frac{x+y}{2} \right) + \sum_{j=2}^n f_j \left(\frac{x-y}{2n-2} \right), f_{n+1}(x) \right) < \varepsilon, \quad (20)$$

for all $x, y \in B$. Setting $x = \frac{x+y}{2}$ and $y = \frac{x-y}{2}$ in (20), we have

$$D \left(f_1 \left(\frac{x}{2} \right) + \sum_{j=2}^n f_j \left(\frac{y}{2n-2} \right), f_{n+1} \left(\frac{x+y}{2} \right) \right) < \varepsilon, \quad (21)$$

for all $x, y \in B$. Letting $x_j = -x$ ($j = 2, \dots, n$) separately, $x_1 = x$ and $x_i = 0$ ($i = 2, \dots, n$ and $i \neq j$) in (17), we get

$$D(f_j(x), f_{n+1}(x)) < \varepsilon, \quad (22)$$

for all $x \in B$, where $j \in \{2, \dots, n\}$. Setting $x = 2x$ and $y = 0$ in (21), we have

$$D(f_1(x), f_{n+1}(x)) < \varepsilon, \quad (23)$$

for all $x \in B$. It follows from (22), (23) and the property (P3) of D that

$$D \left(\sum_{j=1, j \neq k}^n f_j(x), (n-1)f_{n+1}(x) \right) < (n-1)\varepsilon, \quad (24)$$

for all $x \in B$, where $k \in \{1, \dots, n\}$. Putting $x = 2x$ and $y = (2n-2)x$ in (21), we obtain

$$D \left(\sum_{j=1}^n f_j(x), f_{n+1}(nx) \right) < \varepsilon, \quad (25)$$

for all $x \in B$. Using the inequalities (22) to (25) and the properties (P1) to (P3) of D , we conclude that

$$\begin{aligned} & D(f_k(nx), nf_k(x)) \\ &= D(f_k(nx) + f_{n+1}(nx), f_{n+1}(nx) + nf_k(x)) \\ &\leq D(f_k(nx), f_{n+1}(nx)) + D(f_{n+1}(nx), nf_k(x)) \\ &< \varepsilon + D \left(f_{n+1}(nx) + \sum_{j=1, j \neq k}^n f_j(x), \sum_{j=1, j \neq k}^n f_j(x) + f_k(x) + (n-1)f_k(x) \right) \end{aligned}$$

$$\begin{aligned}
&\leq \varepsilon + D\left(f_{n+1}(nx), \sum_{j=1}^n f_j(x)\right) + D\left(\sum_{j=1, j \neq k}^n f_j(x), (n-1)f_k(x)\right) \\
&< \varepsilon + \varepsilon + D\left(\sum_{j=1, j \neq k}^n f_j(x), (n-1)f_k(x)\right) \\
&= 2\varepsilon + D\left(\sum_{j=1, j \neq k}^n f_j(x) + (n-1)f_{n+1}(x), (n-1)f_{n+1}(x) + (n-1)f_k(x)\right) \\
&\leq 2\varepsilon + D\left(\sum_{j=1, j \neq k}^n f_j(x), (n-1)f_{n+1}(x)\right) + D((n-1)f_{n+1}(x), (n-1)f_k(x)) \\
&< 2\varepsilon + (n-1)\varepsilon + D\left(\sum_{j=1, j \neq k}^n f_j(x), (n-1)f_{n+1}(x)\right) \\
&< 2\varepsilon + (n-1)\varepsilon + (n-1)\varepsilon = 2n\varepsilon,
\end{aligned}$$

for all $x \in B$, where $k \in \{1, \dots, n\}$. Rewriting the above, we obtain

$$D\left(f_k(x), \frac{1}{n}f_k(nx)\right) \leq 2\varepsilon,$$

for all $x \in B$, where $k \in \{1, \dots, n\}$. Then, by induction on m , it follows that

$$D\left(f_k(x), \frac{1}{n^m}f_k(n^m x)\right) \leq \sum_{j=0}^{m-1} \frac{2}{n^j}\varepsilon,$$

for all $x \in B$, where $k \in \{1, \dots, n\}$. Using the same method as in the proof of Theorem 3.1, we find a unique additive mapping $\mathcal{G} : X \rightarrow Y$ such that (18) holds for $k \in \{1, \dots, n\}$.

It follows from (23), (24), (25) and the properties (P2) and (P3) of D that

$$\begin{aligned}
&D(f_{n+1}(nx), nf_{n+1}(x)) \\
&= D\left(f_{n+1}(nx) + \sum_{j=1}^n f_j(x), \sum_{j=1}^n f_j(x) + nf_{n+1}(x)\right) \\
&\leq D\left(f_{n+1}(nx), f_1(x) + \sum_{j=2}^n f_j(x)\right) + D\left(f_1(x) + \sum_{j=2}^n f_j(x), nf_{n+1}(x)\right) \\
&\leq D\left(\sum_{j=1}^n f_j(x), f_{n+1}(nx)\right) + D(f_1(x), f_{n+1}(x)) + D\left(\sum_{j=2}^n f_j(x), (n-1)f_{n+1}(x)\right) \\
&< \varepsilon + \varepsilon + (n-1)\varepsilon = (n+1)\varepsilon,
\end{aligned}$$

for all $x \in B$. We deduce from the last inequality that

$$D\left(f_{n+1}(x), \frac{1}{n}f_{n+1}(nx)\right) \leq \frac{n+1}{n}\varepsilon,$$

for all $x \in B$. Then, by induction on m , we get

$$D\left(f_{n+1}(x), \frac{1}{n^m}f_{n+1}(n^m x)\right) \leq \sum_{j=1}^m \frac{n+1}{n^j}\varepsilon,$$

for all $x \in B$. Using the same method as in the proof of Theorem 3.1, we find a unique additive mapping $\mathcal{G} : X \rightarrow Y$ such that (19) holds. \square

Now, we investigate the stability problem for the Pexiderized general linear type fuzzy number-valued functional equation in n variables.

Corollary 4.2. *Let $f_1, \dots, f_{n+1} : B \rightarrow X_F$ be fuzzy number-valued mappings with $f_1(0) = \dots = f_n(0) = \tilde{0}$ such that*

$$D\left(\sum_{i=1}^n A_i f_i(x_i), f_{n+1}\left(\sum_{i=1}^n a_i x_i\right)\right) < \varepsilon, \quad (26)$$

for all $x_1, \dots, x_n \in B$, where $\varepsilon, a_i, A_i > 0$. Then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that

$$D(f_k(x), \mathcal{G}(x)) \leq \frac{2n}{A_k(n-1)}\varepsilon, \quad (k = 1, 2, \dots, n), \quad (27)$$

$$D(f_{n+1}(x), \mathcal{G}(x)) \leq \frac{n+1}{n-1}\varepsilon, \quad (28)$$

for all $x \in B$.

Proof. Setting $x_i = \frac{x_i}{a_i}$ for $i = 1, \dots, n$ in (26), we get

$$D\left(\sum_{i=1}^n A_i f_i\left(\frac{x_i}{a_i}\right), f_{n+1}\left(\sum_{i=1}^n x_i\right)\right) < \varepsilon, \quad (29)$$

for all $x_1, \dots, x_n \in B$. For $i = 1, \dots, n$, define

$$G_i(x_i) := A_i f_i\left(\frac{x_i}{a_i}\right), \quad (30)$$

for all $x_i \in B$. Then, by (29),

$$D\left(\sum_{i=1}^n G_i(x_i), f_{n+1}\left(\sum_{i=1}^n x_i\right)\right) < \varepsilon, \quad (31)$$

for all $x_1, \dots, x_n \in B$. Letting $x_1 = \frac{\sum_{i=1}^n x_i}{n}$ and $x_i = \frac{\sum_{j=1, j \neq i}^n x_j - (n-1)x_i}{n}$ for $i = 2, \dots, n$ in (31), we get

$$D\left(G_1\left(\frac{\sum_{i=1}^n x_i}{n}\right) + \sum_{i=2}^n G_i\left(\frac{\sum_{j=1, j \neq i}^n x_j - (n-1)x_i}{n}\right), f_{n+1}(x_1)\right) < \varepsilon,$$

for all $x_1, \dots, x_n \in B$. Using the same method as in the proofs of Theorems 3.1 and 4.1, the limits

$$\lim_{m \rightarrow \infty} \frac{1}{n^m} G_1(n^m x) = \dots = \lim_{m \rightarrow \infty} \frac{1}{n^m} G_n(n^m x) = \lim_{m \rightarrow \infty} \frac{1}{n^m} f_{n+1}(n^m x),$$

exist for all $x \in B$ and are unique additive mappings satisfying

$$D\left(G_k(x), \lim_{m \rightarrow \infty} \frac{1}{n^m} G_k(n^m x)\right) \leq \frac{2n}{n-1}\varepsilon, \quad (32)$$

$$D\left(f_{n+1}(x), \lim_{m \rightarrow \infty} \frac{1}{n^m} f_{n+1}(n^m x)\right) \leq \frac{n+1}{n-1}\varepsilon, \quad (33)$$

for all $x \in B$, where $k \in \{1, \dots, n\}$. Now, (27) and (28) follow from (30), (32), (33) and the property (P1) of D . \square

Corollary 4.2 with $n = 2$ implies the following.

Corollary 4.3. (Compare with [39, Theorem 2]). *If fuzzy number-valued mappings $f, g, h : B \rightarrow X_F$ with $f(0) = g(0) = \tilde{0}$ satisfy the inequality*

$$D(Af(x) + Bg(y), h(ax + by)) < \varepsilon, \quad (34)$$

for all $x, y \in B$, where $\varepsilon, a, b, A, B > 0$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that

$$D(f(x), \mathcal{G}(x)) \leq \frac{4}{A}\varepsilon, \quad (35)$$

$$D(g(x), \mathcal{G}(x)) \leq \frac{4}{B}\varepsilon, \quad (36)$$

$$D(h(x), \mathcal{G}(x)) \leq 3\varepsilon, \quad (37)$$

for all $x \in B$.

Notation 4.4. Let $f, g, h : B \rightarrow X_F$ be fuzzy number-valued mappings with $f(0) = g(0) = \tilde{0}$. Then

(i) If f, g, h satisfy the inequality (34) with $A = B = a = b = 1$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that (35) to (37) hold for $A = B = 1$.

(ii) If f, g, h satisfy the inequality (34) with $A = B = a = b = \frac{1}{2}$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that (35) to (37) hold for $A = B = \frac{1}{2}$.

(iii) If f, g, h satisfy the inequality (34) with $A = a = t \in (0, 1)$ and $B = b = 1 - t$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that (35) to (37) hold for $A = t$ and $B = 1 - t$.

(iv) If f, g, h satisfy the inequality (34) with $A = a$ and $B = b$, then there exists a unique additive mapping $\mathcal{G} : B \rightarrow X_F$ such that (35) to (37) hold for $A = a$ and $B = b$.

5 Conclusions

In this paper, we proved the stability of several types of additive fuzzy number-valued functional equations, including Cauchy, Jensen, Cauchy-Jensen, Pexiderized Cauchy, Pexiderized Jensen, Pexiderized affine, Pexiderized linear and Pexiderized general linear type fuzzy number-valued functional equations. Our results generalized certain important results obtained by other authors for these equations when they are a single-valued or a set-valued one. Obviously, this paper provided us a novel idea to discuss the stability of functional equations from a more unified perspective. Certainly, further work will focus on the stability of other types of (functional, difference, differential, integral) equations by using this idea.

References

- [1] T. V. An, H. Vu, N. V. Hoa, *Hadamard-type fractional calculus for fuzzy functions and existence theory for fuzzy fractional functional integro-differential equations*, Journal of Intelligent and Fuzzy Systems, **36** (2019), 3591-3605.
- [2] G. A. Anastassiou, S. G. Gal, *On a fuzzy trigonometric approximation theorem of Weierstrass-type*, The Journal of Fuzzy Mathematics, **9** (2001), 701-708.
- [3] J. Ban, *Ergodic theorems for random compact sets and fuzzy variables in Banach spaces*, Fuzzy Sets and Systems, **44** (1991), 71-82.
- [4] B. Bede, S. G. Gal, *Fuzzy-number-valued almost periodic functions*, Fuzzy Sets and Systems, **147** (2004), 385-404.
- [5] A. Bodaghi, Th. M. Rassias, A. Zivari-Kazempour, *A fixed point approach to the stability of additive-quadratic-quartic functional equations*, International Journal of Nonlinear Analysis and Applications, **11** (2020), 17-28.
- [6] J. Brzdęk, A. Pietrzyk, *A note on stability of the general linear equation*, Aequationes Mathematicae, **75** (2008), 267-270.
- [7] J. Brzdęk, D. Popa, B. Xu, *Selections of set-valued maps satisfying a linear inclusion in a single variable*, Nonlinear Analysis: Theory, Methods and Applications, **74** (2011), 324-330.
- [8] T. Cardinali, K. Nikodem, F. Papalini, *Some results on stability and characterization of K -convexity of set-valued functions*, Annales Polonici Mathematici, **58** (1993), 185-192.
- [9] R. Chaharpashlou, A. Atangana, R. Saadati, *On the fuzzy stability results for fractional stochastic Volterra integral equation*, Discrete and Continuous Dynamical Systems-Series S, **14**(10) (2021), 3529-3539.
- [10] R. Chaharpashlou, R. Saadati, *Best approximation of a nonlinear fractional Volterra integro-differential equation in matrix MB-space*, Advances in Difference Equations, **2021** (2021), Doi: 10.1186/s13662-021-03275-2.

- [11] R. Chaharpashlou, R. Saadati, A. Atangana, *Ulam-Hyers-Rassias stability for nonlinear Ψ -Hilfer stochastic fractional differential equation with uncertainty*, Advances in Difference Equations, **2020** (2020), Doi:10.1186/s13662-020-02797-5.
- [12] C. K. Choi, *Stability of Pexiderized Jensen and Jensen type functional equations on restricted domains*, Bulletin of the Korean Mathematical Society, **56** (2019), 801-813.
- [13] H. Y. Chu, A. Kim, S. K. Yoo, *On the stability of the generalized cubic set-valued functional equation*, Applied Mathematics Letters, **37** (2014), 7-14.
- [14] J. Chung, *Hyers-Ulam stability theorems for Pexider equations in the space of Schwartz distributions*, Archiv der Mathematik, **84** (2005), 527-537.
- [15] Z. Gajda, R. Ger, *Subadditive multifunctions and Hyers-Ulam stability*, International Series of Numerical Mathematics, **80** (1987), 281-291.
- [16] A. Ebadian, I. Nikoufar, Th. M. Rassias, N. Ghobadipour, *Stability of generalized derivations on Hilbert C^* -modules associated with a Pexiderized Cauchy-Jensen type functional equation*, Acta Mathematica Scientia, **32** (2012), 1226-1238.
- [17] M. Eshaghi, H. Khodaei, M. Kamyar, *Stability of Cauchy-Jensen type functional equation in generalized fuzzy normed spaces*, Computer and Mathematics with Applications, **62** (2011), 2950-2960.
- [18] M. Hukuhara, *Intégration des applications mesurables dont la valeur est un compact convexe*, Funkcialaj Ekvacioj-Serio Internacia, **10** (1967), 205-223.
- [19] S. Y. Jang, C. Park, Y. Cho, *Hyers-Ulam stability of a generalized additive set-valued functional equation*, Journal of Inequalities and Applications, **2013** (2013), Doi: 10.1186/1029-242X-2013-101.
- [20] K. W. Jun, H. M. Kim, J. M. Rassias, *Extended Hyers-Ulam stability for Cauchy-Jensen mappings*, Journal of Difference Equations and Applications, **13** (2007), 1139-1153.
- [21] H. Khodaei, *On the stability of additive, quadratic, cubic and quartic set-valued functional equations*, Results in Mathematics, **68** (2015), 1-10.
- [22] Y. H. Lee, K. W. Jun, *A generalization of the Hyers-Ulam-Rassias stability of Pexider equation*, Journal of Mathematical Analysis and Applications, **246** (2000), 627-638.
- [23] G. Lu, C. Park, *Hyers-Ulam stability of additive set-valued functional equations*, Applied Mathematics Letters, **24** (2011), 1312-1316.
- [24] D. Mihet, *The fixed point method for fuzzy stability of the Jensen functional equation*, Fuzzy Sets and Systems, **160** (2009), 1663-1667.
- [25] A. Najati, *Homomorphisms in quasi-Banach algebras associated with a Pexiderized Cauchy-Jensen functional equation*, Acta Mathematica Sinica-English Series, **25** (2009), 1529-1542.
- [26] A. Najati, J. I. Kang, Y. J. Cho, *Local stability of the Pexiderized Cauchy and Jensen's equations in fuzzy spaces*, Journal of Inequalities and Applications, **2011** (2011), Doi: 10.1186/1029-242X-2011-78.
- [27] K. Nikodem, *The stability of the Pexider equation*, Annales Mathematicae Silesianae, **5** (1991), 91-93.
- [28] K. Nikodem, D. Popa, *On selections of general linear inclusions*, Publicationes Mathematicae Debrecen, **75** (2009), 239-249.
- [29] K. Nikodem, S. Wąsowicz, *A sandwich theorem and Hyers-Ulam stability of affine functions*, Aequationes Mathematicae, **49** (1995), 160-164.
- [30] C. Park, D. O'Regan, R. Saadati, *Stability of some set-valued functional equations*, Applied Mathematics Letters, **24** (2011), 1910-1914.
- [31] J. V. Pexider, *Notiz über funktional theoreme*, Monatshefte für Mathematik und Physik, **14** (1903), 293-301.

- [32] T. Phochai, S. Saejung, *Some notes on the Ulam stability of the general linear equation*, Acta Mathematica Hungarica, **158** (2019), 40-52.
- [33] V. Y. Protasov, *Stability of affine approximations on bounded domains*, Springer Optimization and Its Applications: Nonlinear Analysis, **68** (2012), 587-606.
- [34] M. Puri, D. Ralescu, *Differentials of fuzzy functions*, Journal of Mathematical Analysis and Applications, **91** (1983), 552-558.
- [35] W. Ren, Z. Yang, X. Sun, M. Qi, *Hyers-Ulam stability of Hermite fuzzy differential equations and fuzzy Mellin transform*, Journal of Intelligent and Fuzzy Systems, **35** (2018), 3721-3731.
- [36] B. V. Senthil Kumar, H. Dutta, S. Sabarinathan, *Fuzzy approximations of a multiplicative inverse cubic functional equation*, Soft Computing, **24** (2020), 13285-13292.
- [37] W. Smajdor, *Superadditive set-valued functions*, Glasnik Matematički, **21** (1986), 343-348.
- [38] H. Vu, J. M. Rassias, N. V. Hoa, *Ulam-Hyers-Rassias stability for fuzzy fractional integral equations*, Iranian Journal of Fuzzy Systems, **17**(2) (2020), 17-27.
- [39] J. R. Wu, Z. Y. Jin, *A note on Ulam stability of some fuzzy number-valued functional equations*, Fuzzy Sets and Systems, **375** (2019), 191-195.